

Fundamental theorems of Doi-Hopf modules in a nonassociative setting

J.N. ALONSO ÁLVAREZ¹, J.M. FERNÁNDEZ VILABOA², R. GONZÁLEZ RODRÍGUEZ³

¹ Departamento de Matemáticas, Universidad de Vigo, Campus Universitario Lagoas-Marcosende, E-36280 Vigo, Spain (e-mail: jnalonso@uvigo.es)

² Departamento de Álgebra, Universidad de Santiago de Compostela. E-15771 Santiago de Compostela, Spain (e-mail: josemanuel.fernandez@usc.es)

³ Departamento de Matemática Aplicada II, Universidad de Vigo, Campus Universitario Lagoas-Marcosende, E-36310 Vigo, Spain (e-mail: rgon@dma.uvigo.es)

Abstract In this paper we introduce the notion of weak non-associative Doi-Hopf module and give the Fundamental Theorem of Hopf modules in this setting. Also we prove that there exists a categorical equivalence that admits as particular instances the ones constructed in the literature for Hopf algebras, weak Hopf algebras, Hopf quasigroups, and weak Hopf quasigroups.

Keywords. Hopf algebra, Weak Hopf algebra, Hopf quasigroup, Weak Hopf quasigroup, Doi-Hopf module, Fundamental Theorem.

MSC 2010: 18D10, 16T05, 17A30, 20N05.

1. INTRODUCTION

Let $\mathbb{F}$ be a field and $\mathcal{C} = \mathbb{F} - Vect$. Let H be a Hopf algebra in $\mathcal{C}$ and let B be a right H -comodule algebra with coaction $\rho_B : B \rightarrow B \otimes H$, $\rho_B(b) = b_{(0)} \otimes b_{(1)}$, where the unadorned tensor product is the tensor product over $\mathbb{F}$ and for $\rho_B(b)$ we used the Sweedler notation. In [13], Doi introduced the notion of (H, B) -Hopf module, as a generalization of the classical notion of Hopf module, defined by Larson and Sweedler in [21], in the following way: Let M be a right B -module and a right H -comodule. If, for all $m \in M$ and $b \in B$, we write $m.b$ for the action and $\rho_M(m) = m_{[0]} \otimes m_{[1]}$ for the coaction, we will say that M is an (H, B) -Hopf module if the equality

$$\rho_M(m.b) = m_{[0]}.b_{(1)} \otimes m_{[1]}b_{(2)}$$

holds, where $m_{[1]}b_{(2)}$ is the product in H of $m_{[1]}$ and $b_{(2)}$. A morphism between two (H, B) -Hopf modules is an $\mathbb{F}$ -linear map that is B -linear and H -colinear. Hopf modules and morphisms of Hopf modules constitute the category of (H, B) -Hopf modules denoted by $\mathcal{M}_B^H$. If there exists a right H -comodule map $h : H \rightarrow B$ which is an algebra map, and $M^{coH} = \{m \in M \mid \rho_M(m) = m \otimes 1_B\}$, $B^{coH} = \{b \in B \mid \rho_B(b) = b \otimes 1_B\}$ are the subobjects of coinvariants, for any $m \in M^{coH}$ and $b \in B^{coH}$ we have that $m.b \in M^{coH}$ and then M^{coH} is a right B^{coH} -module. Using this property Doi proved in Theorem 3 of [13] that M is isomorphic to $M^{coH} \otimes_{B^{coH}} B$ as (H, B) -Hopf modules. Moreover, if N is a right B^{coH} -module, the tensor product $N \otimes_{B^{coH}} B$, with the action and coaction induced by the product of B and the coproduct of H , is an (H, B) -Hopf module. This construction is functorial and then we have a functor, called the induction functor, $F = - \otimes_{B^{coH}} B : \mathcal{C}_{B^{coH}} \rightarrow \mathcal{M}_B^H$. Also, for all $M \in \mathcal{M}_B^H$, the construction of M^{coH} is functorial and we have a functor of coinvariants $G = (-)^{coH} : \mathcal{M}_B^H \rightarrow \mathcal{C}_{B^{coH}}$ such that $F \dashv G$. Moreover, F and G induce a categorical equivalence between $\mathcal{M}_B^H$ and the category of right B^{coH} -modules. This categorical equivalence was called by Doi and Takeuchi in [14], the strong structure theorem for $\mathcal{M}_B^H$, and, for $B = H$ and $h = id_H$, contains as a particular instance the equivalence derived of the Fundamental Theorem of Hopf modules proved by Larson and Sweedler (see [21], [30]).

The categorical equivalence of the previous paragraph remains valid for weak Hopf algebras. For a weak Hopf algebra H , Böhm introduced in [8] the category of Hopf modules, denoted by $\mathcal{M}_H^H$, in the same way that in the Hopf algebra setting. If $M \in \mathcal{M}_H^H$, the subobject of coinvariants is defined by $M^{coH} = \{m \in M \mid \rho_M(m) = m_{[0]} \otimes \Pi_H^L(m_{[1]})\}$, where Π_H^L is the target morphism associated to H . In [8] we can find the weak version of the Fundamental Theorem of Hopf modules, i.e.: For all Hopf module M , $M^{coH} \otimes_{H_L} H$ is isomorphic to M as Hopf modules, where H_L is the image of Π_H^L . Moreover, if $\mathcal{C}_{H_L}$ is the category of right H_L -modules, there exist two functors $F = - \otimes_{H_L} H : \mathcal{C}_{H_L} \rightarrow \mathcal{M}_H^H$ and $G = ()^{coH} : \mathcal{M}_H^H \rightarrow \mathcal{C}_{H_L}$ such that F is left adjoint of G and they induce a pair of inverse equivalences. Therefore, in the weak setting, $\mathcal{M}_H^H$ is equivalent to $\mathcal{C}_{H_L}$. In this case, the following property is a relevant fact for subsequent generalizations: there is an isomorphism of Hopf modules between the tensor product $M^{coH} \otimes_{H_L} H$ and $M^{coH} \times H$, where $M^{coH} \times H$ is the image of a suitable idempotent morphism $\nabla_M : M^{coH} \otimes H \rightarrow M^{coH} \otimes H$. Later, in [9], Böhm introduced the notion of weak Doi-Hopf module (or weak (H, B) -Doi-Hopf module), associated to a weak Hopf algebra H and a right H -comodule algebra B , and the category of weak Doi-Hopf modules denoted as in the non-weak setting by $\mathcal{M}_B^H$. In 2004, Zhang and Zhu proved that for any weak Doi-Hopf module M (also called by these authors weak (H, B) -Doi-Hopf module), if there exists a right H -comodule map $h : H \rightarrow B$ which is an algebra map, the objects M and $M^{coH} \otimes_{B^{coH}} B$ are isomorphic as (H, B) -Doi-Hopf modules. In this case $B^{coH} = \{b \in B \mid \rho_B(b) = b_{(0)} \otimes \Pi_H^L(b_{(1)})\}$ and, if $B = H$ and $h = id_H$, they recover the isomorphism constructed by Böhm in [8]. As in the Hopf setting, it is possible to construct the induction functor $F = - \otimes_{B^{coH}} B : \mathcal{C}_{B^{coH}} \rightarrow \mathcal{M}_B^H$ and the functor of coinvariants $G = ()^{coH} : \mathcal{M}_B^H \rightarrow \mathcal{C}_{B^{coH}}$. These functors satisfy that $F \dashv G$ and F and G is a pair of inverse equivalences. Therefore, $\mathcal{M}_B^H$ is equivalent to the category of right B^{coH} -modules (see [16]).

In the two previous paragraphs we wrote about categorical equivalences for categories of Hopf modules connected to associative algebraic structures like Hopf algebras and weak Hopf algebras. An interesting generalization of Hopf algebras are nonassociative Hopf algebras. As in the quasi-Hopf setting, nonassociative Hopf algebras are not associative, but the lack of this property is compensated in this case by some axioms involving the division operation. The notion of nonassociative Hopf algebra in a category of vector spaces was introduced by Pérez- Izquierdo [25] with the aim of to construct the universal enveloping algebra for Sabinin algebras, prove a Poincaré-Birkhoff-Witt Theorem for Sabinin algebras and give a nonassociative version of the Milnor-Moore theorem. Later, Klim and Majid [20], in order to understand the structure and relevant properties of the algebraic 7-sphere, introduced the notion of Hopf quasigroup. Hopf quasigroups are examples of nonassociative Hopf algebras and in recent years, interesting research about its specific structure and its dual has been developed ([10], [17], [11], [12], [18], [19], [15], [1], [2]). Moreover, nonassociative Hopf algebras arise naturally related with other structures in various nonassociative contexts like, for example quantum quasigroups in the sense of Smith ([26], [27], [28], [29]). Nonassociative Hopf algebras include the example of an enveloping algebra $U(L)$ of a Malcev algebra (see [24], [20], [31]) as well as the notion of the loop algebra RL of a loop L (see [7], [23]). Then, nonassociative Hopf algebras unify Moufang loops and Malcev algebras, and, more generally, formal loops and Sabinin algebras, in the same way that Hopf algebras unify groups and Lie algebras.

For a Hopf quasigroup in the sense of [20], Brzeziński defined in [10] the notion of Hopf module obtaining a categorical equivalence as in the associative context. In this case, the main difference appears in the definition of the category of Hopf modules $\mathcal{M}_H^H$. Firstly, because the notion of Hopf module reflects the non-associativity of the product defined on H . Secondly, the morphisms are H -quasilinear and H -colinear (see Definition 3.4 of [10]). In Lemma 3.5 of [10], we can find that, if $M \in \mathcal{M}_H^H$ and M^{coH} is defined like in the Hopf algebra setting, M is isomorphic to $M^{coH} \otimes H$ as Hopf modules. Therefore the Fundamental Theorem of Hopf modules also holds for Hopf quasigroups. Moreover, there exist two functors $F = - \otimes H : \mathcal{C} \rightarrow \mathcal{M}_H^H$ and $G = ()^{coH} : \mathcal{M}_H^H \rightarrow \mathcal{C}$ such that $F \dashv G$, and they induce a pair of inverse equivalences. Thus, as it occurs in the Hopf algebra ambit, $\mathcal{M}_H^H$ is equivalent to the category of $\mathbb{F}$ -vector spaces.

Hopf quasigroups admit a generalization to the weak setting. The new notion, called weak Hopf quasigroup, was introduced in [3] in a monoidal context and a family of non trivial examples can be

obtained by working with bigroupoids, i.e., bicategories where every 1-cell is an equivalence and every 2-cell is an isomorphism (see Example 2.3 of [3]). In [5] we described these algebraic objects in terms of fusion morphisms and in [3], for a weak Hopf quasigroup H in a braided monoidal category $\mathcal{C}$ with tensor product $\otimes$, using the ideas proposed by Brzeziński for Hopf quasigroups, we introduce the notion of Hopf module and the category of Hopf modules $\mathcal{M}_H^H$. In this case, if we define M^{coH} in the same way that in the weak Hopf algebra setting, we obtain the weak nonassociative version of the Fundamental Theorem of Hopf modules in the following way: every Hopf module M is isomorphic to $M^{coH} \times H$ as Hopf modules, where $M^{coH} \times H$ is the image of the same idempotent ∇_M used for Hopf modules associated to a weak Hopf algebra. Moreover, in [4] we proved that H_L , the image of the target morphism, is a monoid, and then it is possible to take into consideration the category $\mathcal{C}_{H_L}$ to construct the tensor product $M^{coH} \otimes_{H_L} H$, and, if the functor $-\otimes H$ preserves coequalizers, to endow this object with a Hopf module structure. Unfortunately, unlike the case of weak Hopf algebras, it is not possible to assure in general that $M^{coH} \otimes_{H_L} H$ is isomorphic to $M^{coH} \times H$. In order to find sufficient conditions under which these objects are isomorphic in $\mathcal{M}_H^H$, we introduce in [6] the category of strong Hopf modules, denoted by $\mathcal{SM}_H^H$ and obtain that there exist two functors $F = -\otimes_{H_L} H : \mathcal{C}_{H_L} \rightarrow \mathcal{SM}_H^H$ and $G = (-)^{coH} : \mathcal{SM}_H^H \rightarrow \mathcal{C}_{H_L}$ such that F is left adjoint of G and they induce a pair of inverse equivalences. In the Hopf quasigroup setting every Hopf module is strong, and then our results are the ones proved by Brzeziński in [10]. The same happens in the weak Hopf case and then we generalize the theorem proved by Böhm, Nill and Szlachányi in [8].

Let $\mathcal{C}$ be a braided monoidal category with tensor product $\otimes$. Then for a weak Hopf quasigroup H in $\mathcal{C}$ and a right H -comodule magma B (see [4] for the definition), a question naturally arises: Is it possible to define a general category of (H, B) -Hopf modules and to prove a general theorem that permit to recover as particular instances the categorical equivalences cited in the previous paragraphs? The main contribution of this paper is to give a positive answer to this question.

Now, we describe the paper in detail. After this introduction, for a weak Hopf quasigroup H and a right H -comodule magma B in a strict braided monoidal category $\mathcal{C}$ where every idempotent morphism splits, in the second section we introduce the notion of anchor morphism $h : H \rightarrow B$ as an H -comodule morphism such that it is a morphism of unital magmas satisfying two suitable conditions. For an anchor morphism h , in Definition 2.8, we define the notion of strong (H, B, h) -Hopf module and prove some properties of these modules. We also find the condition under which the subobject of coinvariants of B , defined as in the weak Hopf algebra context, i.e., $B^{coH} = \{b \in B \mid \rho_B(b) = b_{(0)} \otimes \Pi_H^L(b_{(1)})\}$, is a monoid, and construct the new category of strong (H, B, h) -Hopf modules, denoted by $\mathcal{SM}_B^H$. Moreover, if the category $\mathcal{C}$ admits coequalizers and the functors $-\otimes B$ and $-\otimes H$ preserve coequalizers, we prove in Theorem 2.22 that the Fundamental Theorem of Hopf Modules holds. In other words, for any strong (H, B, h) -Hopf module the objects M and $M^{coH} \otimes_{B^{coH}} B$ are isomorphic in $\mathcal{SM}_B^H$. This result admits as particular instances the results with the same name cited in the previous paragraphs for associative and nonassociative (weak) Hopf structures. Finally, in the last section, we define the induction functor $F = -\otimes_{B^{coH}} B : \mathcal{C}_{B^{coH}} \rightarrow \mathcal{SM}_B^H$ and the functor of coinvariants $G = (-)^{coH} : \mathcal{SM}_B^H \rightarrow \mathcal{C}_{B^{coH}}$, proving that $F \dashv G$. Also, F and G is a pair of inverse equivalences and, therefore, $\mathcal{SM}_B^H$ is equivalent to the category of right B^{coH} -modules.

Throughout this paper $\mathcal{C}$ denotes a strict braided monoidal category with tensor product $\otimes$, unit object K and braid c . For each object M in $\mathcal{C}$, we denote the identity morphism by $id_M : M \rightarrow M$ and, for simplicity of notation, given objects M , N and P in $\mathcal{C}$ and a morphism $f : M \rightarrow N$, we write $P \otimes f$ for $id_P \otimes f$ and $f \otimes P$ for $f \otimes id_P$. We also assume that every idempotent morphism in $\mathcal{C}$ splits, i.e., if $\nabla : Y \rightarrow Y$ is such that $\nabla = \nabla \circ \nabla$, there exist an object Z , called the image of p , and morphisms $i : Z \rightarrow Y$ and $p : Y \rightarrow Z$ such that $\nabla = i \circ p$ and $p \circ i = id_Z$. The morphisms p and i will be called a factorization of q . Note that Z , p and i are unique up to isomorphism. The categories satisfying this property constitute a broad class that includes, among others, the categories with epi-monic decomposition for morphisms and categories with equalizers or coequalizers.

As for prerequisites, the reader is expected to be familiar with the notions of (co)unital (co)magma, (co)monoid, and morphism of (co)unital (co)magmas. By a unital magma in $\mathcal{C}$ we understand a triple $A = (A, \eta_A, \mu_A)$ where A is an object in $\mathcal{C}$ and $\eta_A : K \rightarrow A$ (unit), $\mu_A : A \otimes A \rightarrow A$ (product) are morphisms in $\mathcal{C}$ such that $\mu_A \circ (A \otimes \eta_A) = id_A = \mu_A \circ (\eta_A \otimes A)$. If μ_A is associative, that is, $\mu_A \circ (A \otimes \mu_A) = \mu_A \circ (\mu_A \otimes A)$, the unital magma will be called a monoid in $\mathcal{C}$. Given two unital magmas (monoids) $A = (A, \eta_A, \mu_A)$ and $B = (B, \eta_B, \mu_B)$, $f : A \rightarrow B$ is a morphism of unital magmas (monoids) if $\mu_B \circ (f \otimes f) = f \circ \mu_A$ and $f \circ \eta_A = \eta_B$.

By duality, a counital comagma in $\mathcal{C}$ is a triple $D = (D, \varepsilon_D, \delta_D)$ where D is an object in $\mathcal{C}$ and $\varepsilon_D : D \rightarrow K$ (counit), $\delta_D : D \rightarrow D \otimes D$ (coproduct) are morphisms in $\mathcal{C}$ such that $(\varepsilon_D \otimes D) \circ \delta_D = id_D = (D \otimes \varepsilon_D) \circ \delta_D$. If δ_D is coassociative, that is, $(\delta_D \otimes D) \circ \delta_D = (D \otimes \delta_D) \circ \delta_D$, the counital comagma will be called a comonoid. If $D = (D, \varepsilon_D, \delta_D)$ and $E = (E, \varepsilon_E, \delta_E)$ are counital comagmas (comonoids), $f : D \rightarrow E$ is a morphism of counital comagmas (comonoids) if $(f \otimes f) \circ \delta_D = \delta_E \circ f$ and $\varepsilon_E \circ f = \varepsilon_D$.

If A, B are unital magmas (monoids) in $\mathcal{C}$, the object $A \otimes B$ is a unital magma (monoid) in $\mathcal{C}$ where $\eta_{A \otimes B} = \eta_A \otimes \eta_B$ and $\mu_{A \otimes B} = (\mu_A \otimes \mu_B) \circ (A \otimes c_{B,A} \otimes B)$. In a dual way, if D, E are counital comagmas (comonoids) in $\mathcal{C}$, $D \otimes E$ is a counital comagma (comonoid) in $\mathcal{C}$ where $\varepsilon_{D \otimes E} = \varepsilon_D \otimes \varepsilon_E$ and $\delta_{D \otimes E} = (D \otimes c_{D,E} \otimes E) \circ (\delta_D \otimes \delta_E)$.

Let A be a monoid. The pair (M, ϕ_M) is a right A -module if M is an object in $\mathcal{C}$ and $\phi_M : M \otimes A \rightarrow M$ is a morphism in $\mathcal{C}$ satisfying $\phi_M \circ (M \otimes \eta_A) = id_M$, $\phi_M \circ (\phi_M \otimes A) = \phi_M \circ (M \otimes \mu_A)$. Given two right A -modules (M, ϕ_M) and (N, ϕ_N) , $f : M \rightarrow N$ is a morphism of right A -modules if $\phi_N \circ (f \otimes A) = f \circ \phi_M$. If D is a comonoid, the pair (M, ρ_M) is a right D -comodule if M is an object in $\mathcal{C}$ and $\rho_M : M \rightarrow M \otimes D$ is a morphism in $\mathcal{C}$ satisfying $(M \otimes \varepsilon_D) \circ \rho_M = id_M$, $(\rho_M \otimes D) \circ \rho_M = (M \otimes \delta_D) \circ \rho_M$. Given two right D -comodules (M, ρ_M) and (N, ρ_N) , $f : M \rightarrow N$ is a morphism of right D -comodules if $(f \otimes D) \circ \rho_M = \rho_N \circ f$.

Finally, if D is a comagma and A a magma, given two morphisms $f, g : D \rightarrow A$ we will denote by $f * g$ its convolution product in $\mathcal{C}$, that is

$$f * g = \mu_A \circ (f \otimes g) \circ \delta_D.$$

2. DOI-HOPF MODULES FOR WEAK HOPF QUASIGROUPS

We begin this section by recalling the notion of weak Hopf quasigroup in a braided monoidal category introduced in [3]. In this reference the interested reader can find an exhaustive list of properties of weak Hopf quasigroups, that we will need along the paper.

Definition 2.1. A weak Hopf quasigroup H in $\mathcal{C}$ is a unital magma (H, η_H, μ_H) and a comonoid $(H, \varepsilon_H, \delta_H)$ such that the following axioms hold:

- (a1) $\delta_H \circ \mu_H = (\mu_H \otimes \mu_H) \circ \delta_{H \otimes H}$.
- (a2) $\varepsilon_H \circ \mu_H \circ (\mu_H \otimes H) = \varepsilon_H \circ \mu_H \circ (H \otimes \mu_H)$
 $= ((\varepsilon_H \circ \mu_H) \otimes (\varepsilon_H \circ \mu_H)) \circ (H \otimes \delta_H \otimes H)$
 $= ((\varepsilon_H \circ \mu_H) \otimes (\varepsilon_H \circ \mu_H)) \circ (H \otimes (c_{H,H}^{-1} \circ \delta_H) \otimes H)$.
- (a3) $(\delta_H \otimes H) \circ \delta_H \circ \eta_H = (H \otimes \mu_H \otimes H) \circ ((\delta_H \circ \eta_H) \otimes (\delta_H \circ \eta_H))$
 $= (H \otimes (\mu_H \circ c_{H,H}^{-1}) \otimes H) \circ ((\delta_H \circ \eta_H) \otimes (\delta_H \circ \eta_H))$.
- (a4) There exists $\lambda_H : H \rightarrow H$ in $\mathcal{C}$ (called the antipode of H) such that, if we denote the morphisms $id_H * \lambda_H$ by Π_H^L (target morphism) and $\lambda_H * id_H$ by Π_H^R (source morphism),
 - (a4-1) $\Pi_H^L = ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes c_{H,H}) \circ ((\delta_H \circ \eta_H) \otimes H)$.
 - (a4-2) $\Pi_H^R = (H \otimes (\varepsilon_H \circ \mu_H)) \circ (c_{H,H} \otimes H) \circ (H \otimes (\delta_H \circ \eta_H))$.
 - (a4-3) $\lambda_H * \Pi_H^L = \Pi_H^R * \lambda_H = \lambda_H$.
 - (a4-4) $\mu_H \circ (\lambda_H \otimes \mu_H) \circ (\delta_H \otimes H) = \mu_H \circ (\Pi_H^R \otimes H)$.
 - (a4-5) $\mu_H \circ (H \otimes \mu_H) \circ (H \otimes \lambda_H \otimes H) \circ (\delta_H \otimes H) = \mu_H \circ (\Pi_H^L \otimes H)$.
 - (a4-6) $\mu_H \circ (\mu_H \otimes \lambda_H) \circ (H \otimes \delta_H) = \mu_H \circ (H \otimes \Pi_H^L)$.
 - (a4-7) $\mu_H \circ (\mu_H \otimes H) \circ (H \otimes \lambda_H \otimes H) \circ (H \otimes \delta_H) = \mu_H \circ (H \otimes \Pi_H^R)$.

Note that, if in the previous definition the triple (H, η_H, μ_H) is a monoid, we obtain the notion of weak Hopf algebra in a symmetric monoidal category. Then, if $\mathcal{C}$ is the category of vector spaces over a field

$\mathbb{F}$, we have the original definition of weak Hopf algebra introduced by Böhm, Nill and Szlachányi in [8]. On the other hand, under these conditions, if ε_H and δ_H are morphisms of unital magmas (equivalently, η_H, μ_H are morphisms of counital comagmas), $\Pi_H^L = \Pi_H^R = \eta_H \otimes \varepsilon_H$. As a consequence, conditions (a2), (a3), (a4-1)-(a4-3) trivialize, and we get the notion of Hopf quasigroup defined by Klim and Majid in [20].

By Proposition 3.2 of [3] we know that the antipode is unique, and satisfies that $\lambda_H \circ \eta_H = \eta_H$, $\varepsilon_H \circ \lambda_H = \varepsilon_H$. Also, by Theorem 3.19 of [3], we have that it is antimultiplicative and anticomultiplicative. Moreover, if we define the morphisms $\overline{\Pi}_H^L$ and $\overline{\Pi}_H^R$ by

$$\overline{\Pi}_H^L = (H \otimes (\varepsilon_H \circ \mu_H)) \circ ((\delta_H \circ \eta_H) \otimes H), \quad \overline{\Pi}_H^R = ((\varepsilon_H \circ \mu_H) \otimes H) \circ (H \otimes (\delta_H \circ \eta_H)),$$

we proved in Proposition 3.4 of [3], that $\Pi_H^L, \Pi_H^R, \overline{\Pi}_H^L$ and $\overline{\Pi}_H^R$ are idempotent.

Lemma 2.2. *Let H be a weak Hopf quasigroup and $\Pi \in \{\Pi_H^L, \Pi_H^R, \overline{\Pi}_H^L, \overline{\Pi}_H^R\}$. The following identities hold:*

$$\mu_H \circ ((\mu_H \circ (\Pi \otimes H)) \otimes H) = \mu_H \circ (\Pi \otimes \mu_H), \quad (1)$$

$$\mu_H \circ (H \otimes (\mu_H \circ (\Pi \otimes H))) = \mu_H \circ ((\mu_H \circ (H \otimes \Pi)) \otimes H), \quad (2)$$

$$\mu_H \circ (H \otimes (\mu_H \circ (H \otimes \Pi))) = \mu_H \circ (\mu_H \otimes \Pi). \quad (3)$$

Proof. The proof for Π_H^L is in Proposition 2.4 of [4] and in a similar way we can prove the result for Π_H^R (see also Proposition 2.3 of [5]). The equalities for $\overline{\Pi}_H^L$ and $\overline{\Pi}_H^R$ follow from Proposition 3.11 of [3]. $\square$

If H_L is the image of the idempotent morphism Π_H^L , and $p_L : H \rightarrow H_L, i_L : H_L \rightarrow H$ are the morphisms such that $\Pi_H^L = i_L \circ p_L$ and $p_L \circ i_L = id_{H_L}$, by Proposition 3.13 of [3], i_L is the equalizer of δ_H and $(H \otimes \Pi_H^L) \circ \delta_H$ and p_L is the coequalizer of μ_H and $\mu_H \circ (H \otimes \Pi_H^L)$. Then the triple $(H_L, \varepsilon_{H_L} = \varepsilon_H \circ i_L, \delta_H = (p_L \otimes p_L) \circ \delta_H \circ i_L)$ is a comonoid in $\mathcal{C}$, and as a consequence of Lemma 2.2, $(H_L, \eta_{H_L} = p_L \circ \eta_H, \mu_{H_L} = p_L \circ \mu_H \circ (i_L \otimes i_L))$ is a monoid in $\mathcal{C}$. Following Remark 3.15 of [3], we have similar results for the image of the idempotent morphism Π_H^R denoted by H_R .

Definition 2.3. Let H be a weak Hopf quasigroup and let B be a unital magma, which is also a right H -comodule with coaction $\rho_B : B \rightarrow B \otimes H$ such that

$$\mu_{B \otimes H} \circ (\rho_B \otimes \rho_B) = \rho_B \circ \mu_B. \quad (4)$$

We will say that (B, ρ_B) is a right H -comodule magma if any of the following equivalent conditions hold:

- (b1) $(\rho_B \otimes H) \circ \rho_B \circ \eta_B = (B \otimes (\mu_H \circ c_{H,H}) \otimes H) \circ ((\rho_B \circ \eta_B) \otimes (\delta_H \circ \eta_H))$.
- (b2) $(\rho_B \otimes H) \circ \rho_B \circ \eta_B = (B \otimes \mu_H \otimes H) \circ ((\rho_B \circ \eta_B) \otimes (\delta_H \circ \eta_H))$.
- (b3) $(B \otimes \overline{\Pi}_H^R) \circ \rho_B = (\mu_B \otimes H) \circ (B \otimes (\rho_B \circ \eta_B))$,
- (b4) $(B \otimes \Pi_H^L) \circ \rho_B = ((\mu_B \circ c_{B,B}) \otimes H) \circ (B \otimes (\rho_B \circ \eta_B))$.
- (b5) $(B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ \eta_B = \rho_B \circ \eta_B$.
- (b6) $(B \otimes \Pi_H^L) \circ \rho_B \circ \eta_B = \rho_B \circ \eta_B$.

This definition is similar to the notion of right H -comodule monoid in the weak Hopf algebra setting and the proof for the equivalence of (b1)-(b6) is the same.

Note that, if H is a Hopf quasigroup and B is a unital magma which is also a right H -comodule with coaction $\rho_B : B \rightarrow B \otimes H$, we will say that (B, ρ_B) is a right H -comodule magma if it satisfies (4) and $\rho_B \circ \eta_B = \eta_H \otimes \eta_B$. In this case (b1)-(b6) trivialize.

For example, if H is a (weak) Hopf quasigroup, (H, δ_H) is a right H -comodule magma.

Definition 2.4. Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. We will say that $h : H \rightarrow B$ is an integral if it is a morphism of right H -comodules. The integral will be called total if $h \circ \eta_H = \eta_B$.

Proposition 2.5. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be a total integral. The endomorphism $q_B := \mu_B \circ (B \otimes (h \circ \lambda_H)) \circ \rho_B : B \rightarrow B$ satisfies*

$$\rho_B \circ q_B = (B \otimes \Pi_H^L) \circ \rho_B \circ q_B, \quad (5)$$

$$\rho_B \circ q_B = (B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ q_B, \quad (6)$$

and, as a consequence, q_B is an idempotent morphism. Moreover, if B^{coH} (object of coinvariants) is the image of q_B and $p_B : B \rightarrow B^{coH}$, $i_B : B^{coH} \rightarrow B$ are the morphisms such that $q_B = i_B \circ p_B$ and $id_{B^{coH}} = p_B \circ i_B$,

$$\begin{array}{ccc} B^{coH} & \xrightarrow{i_B} & B \\ & & \downarrow \rho_B \\ & & B \otimes H \\ & & \uparrow (B \otimes \overline{\Pi}_H^R) \circ \rho_B \end{array}$$

$$\begin{array}{ccc} B^{coH} & \xrightarrow{i_B} & B \\ & & \downarrow \rho_B \\ & & B \otimes H \\ & & \uparrow (B \otimes \Pi_H^L) \circ \rho_B \end{array}$$

are equalizer diagrams.

Proof. First note that, by the naturality of c , the condition of right H -comodule morphism for h , and the equality $h \circ \eta_H = \eta_B$, we obtain that

$$h \circ \Pi_H^R = (B \otimes (\varepsilon_H \circ \mu_H)) \circ (c_{H,B} \otimes H) \circ (H \otimes (\rho_B \circ \eta_B)) \quad (7)$$

holds. Then, as a consequence, by (4) and the properties of ε_H and η_B , we have

$$\mu_B \circ (B \otimes (h \circ \Pi_H^R)) \circ \rho_B = id_B. \quad (8)$$

Also,

$$\begin{aligned} \rho_B \circ q_B &= \mu_{B \otimes H} \circ (\rho_B \otimes (\rho_B \circ h \circ \lambda_H)) \circ \rho_B \quad (4) \\ &= (\mu_B \otimes H) \circ (B \otimes ((h \otimes \mu_H) \circ (c_{H,H} \otimes H) \circ (H \otimes (\delta_H \circ \lambda_H)) \circ \delta_H)) \circ \rho_B \quad (\text{comodule condition for } B \text{ and} \\ &\quad \text{comodule morphism condition for } h) \\ &= (\mu_B \otimes H) \circ (B \otimes ((h \otimes \mu_H) \circ (c_{H,H} \otimes H) \circ (H \otimes c_{H,H}) \circ (((H \otimes \lambda_H) \circ \delta_H) \otimes \lambda_H) \circ \delta_H)) \circ \rho_B \\ &\quad (\text{antimultiplicativity of } \lambda_H \text{ and coassociativity of } \delta_H) \\ &= ((\mu_B \circ (B \otimes h)) \otimes \Pi_H^L) \circ (B \otimes (c_{H,H} \circ (H \otimes \lambda_H) \circ \delta_H)) \circ \rho_B \quad (\text{naturality of } c), \end{aligned}$$

and then using that Π_H^L is an idempotent morphism we obtain (5). Therefore, by (34) of [3], we have (6) because

$$(B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ q_B = (B \otimes (\overline{\Pi}_H^R \circ \Pi_H^L)) \circ \rho_B \circ q_B = (B \otimes \Pi_H^L) \circ \rho_B \circ q_B = \rho_B \circ q_B.$$

Then, q_B is an idempotent morphism. Indeed,

$$\begin{aligned} q_B \circ q_B &= \mu_H \circ (B \otimes (h \circ \lambda_H \circ \overline{\Pi}_H^R)) \circ \rho_B \circ q_B \quad (6) \\ &= \mu_B \circ (B \otimes (h \circ \Pi_H^R)) \circ \rho_B \circ q_B \quad ((40) \text{ of } [3]) \\ &= q_B \quad (8). \end{aligned}$$

On the other hand, by (40) of [3] and (8),

$$\mu_B \circ (B \otimes (h \circ \lambda_H \circ \overline{\Pi}_H^R)) \circ \rho_B = \mu_B \circ (B \otimes (h \circ \Pi_H^R)) \circ \rho_B = id_B.$$

Then,

$$\begin{array}{ccc} B^{coH} & \xrightarrow{i_B} & B \\ & & \downarrow \rho_B \\ & & B \otimes H \\ & & \uparrow (B \otimes \overline{\Pi}_H^R) \circ \rho_B \end{array}$$

is a split cofork (see [22]) and thus an equalizer diagram. As a consequence, by (33) of [3], we have

$$(B \otimes \Pi_H^L) \circ \rho_B \circ q_B = (B \otimes (\Pi_H^L \circ \overline{\Pi}_H^R)) \circ \rho_B \circ q_B = (B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ q_B = \rho_B \circ q_B$$

and q_B equalizes ρ_B and $(B \otimes \Pi_H^L) \circ \rho_B$. Therefore, i_B equalizes ρ_B and $(B \otimes \Pi_H^L) \circ \rho_B$ because $p_B \circ i_B = id_{B^{coH}}$. Moreover, if $t : D \rightarrow B$ is such that $(B \otimes \Pi_H^L) \circ \rho_B \circ t = \rho_B \circ t$, composing with $B \otimes \overline{\Pi}_H^R$, we obtain that $(B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ t = \rho_B \circ t$. Thus, there exists a unique morphism $t' : D \rightarrow B^{coH}$ such that $i_B \circ t' = t$. Then,

$$B^{coH} \xrightarrow{i_B} B \xrightleftharpoons[(B \otimes \Pi_H^L) \circ \rho_B]{\rho_B} B \otimes H$$

is an equalizer diagram. $\square$

Remark 2.6. Note that, under the conditions of the previous proposition, the object of coinvariants is independent of the total integral h because it is the equalizer object of ρ_B and $(B \otimes \overline{\Pi}_H^R) \circ \rho_B$ (or equivalently, ρ_B and $(B \otimes \Pi_H^L) \circ \rho_B$).

Moreover, (see [4]) the triple $(B^{coH}, \eta_{B^{coH}}, \mu_{B^{coH}})$ is a unital magma (the submagma of coinvariants of B), where $\eta_{B^{coH}} : K \rightarrow B^{coH}$, $\mu_{B^{coH}} : B^{coH} \otimes B^{coH} \rightarrow B^{coH}$ are the factorizations through i_B of the morphisms η_B and $\mu_B \circ (i_B \otimes i_B)$, respectively. Therefore, $\eta_{B^{coH}}$ is the unique morphism such that

$$\eta_B = i_B \circ \eta_{B^{coH}}, \quad (9)$$

and $\mu_{B^{coH}}$ is the unique morphism satisfying

$$\mu_B \circ (i_B \otimes i_B) = i_B \circ \mu_{B^{coH}}. \quad (10)$$

Thus,

$$\eta_{B^{coH}} = p_B \circ \eta_B, \quad (11)$$

and

$$\mu_{B^{coH}} = p_B \circ \mu_B \circ (i_B \otimes i_B). \quad (12)$$

In what follows, the object of coinvariants B^{coH} will be called the submagma of coinvariants of B . Note that, if $B = H$ and $\rho_B = \delta_H$, the submagma of coinvariants is $H^{coH} = H_L$ and then, in this case, it is a monoid.

Definition 2.7. Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. We will say that $h : H \rightarrow B$ is an anchor morphism if it is a multiplicative total integral (i.e., a right H -comodule morphism such that it is a morphism of unital magmas) and the following equalities hold:

- (c1) $\mu_B \circ ((\mu_B \circ (B \otimes h)) \otimes (h \circ \lambda_H)) \circ (B \otimes \delta_H) = \mu_B \circ (B \otimes (h \circ \Pi_H^L))$.
- (c2) $\mu_B \circ ((\mu_B \circ (B \otimes (h \circ \lambda_H))) \otimes h) \circ (B \otimes \delta_H) = \mu_B \circ (B \otimes (h \circ \Pi_H^R))$.

Note that, if the product on B is associative, every multiplicative total integral h satisfies (c1)-(c2) and therefore is an anchor morphism. Also, using that h is a comodule morphism, the condition (c1) can be rewritten as

$$\mu_B \circ (\mu_B \otimes (h \circ \lambda_H)) \circ (B \otimes (\rho_B \circ h)) = \mu_B \circ (B \otimes (h \circ \Pi_H^L)). \quad (13)$$

Moreover, by the definition of weak Hopf quasigroup and (40) of [3], the identity morphism id_H is an anchor morphism for the right H -comodule magma (H, δ_H) .

On the other hand,

$$(B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ \mu_B = (\mu_B \otimes \overline{\Pi}_H^R) \circ (B \otimes \rho_B). \quad (14)$$

Indeed:

$$\begin{aligned} & (B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ \mu_B \\ &= (\mu_B \otimes (\varepsilon_H \circ \mu_H \circ (\mu_H \otimes H))) \otimes H \circ (B \otimes c_{H,B} \otimes H \otimes (\delta_H \circ \eta_H)) \circ (\rho_B \otimes \rho_B) \text{ ((4) and definition of } \overline{\Pi}_H^R) \\ &= (\mu_B \otimes (((\varepsilon_H \circ \mu_H) \otimes (\varepsilon_H \circ \mu_H)) \circ (H \otimes \delta_H \otimes H))) \otimes H \circ (B \otimes c_{H,B} \otimes H \otimes (\delta_H \circ \eta_H)) \circ (\rho_B \otimes \rho_B) \\ & \quad \text{(a2) of Definition 2.1} \\ &= (B \otimes \varepsilon_H \otimes \varepsilon_H \otimes H) \circ (\mu_{B \otimes H} \otimes \mu_H \otimes H) \circ (\rho_B \otimes \rho_B \otimes H \otimes (\delta_H \circ \eta_H)) \circ (B \otimes \rho_B) \text{ (comodule condition for } B) \end{aligned}$$

$$\begin{aligned}
&= (((B \otimes \varepsilon_H) \circ \rho_B \circ \mu_B) \otimes \overline{\Pi}_H^R) \circ (B \otimes \rho_B) \text{ ((4) and definition of } \overline{\Pi}_H^R) \\
&= (\mu_B \otimes \overline{\Pi}_H^R) \circ (B \otimes \rho_B) \text{ (comodule condition for } B).
\end{aligned}$$

As a consequence, if $h : H \rightarrow B$ is a total integral, the equality

$$\mu_B \circ (\mu_B \otimes (h \circ \Pi_H^R)) \circ (B \otimes \rho_B) = \mu_B \quad (15)$$

holds because

$$\begin{aligned}
&\mu_B \circ (\mu_B \otimes (h \circ \Pi_H^R)) \circ (B \otimes \rho_B) \\
&= \mu_B \circ (\mu_B \otimes (h \circ \lambda_H \circ \overline{\Pi}_H^R)) \circ (B \otimes \rho_B) \text{ ((40) of [3])} \\
&= \mu_B \circ (\mu_B \otimes (h \circ \lambda_H \circ \overline{\Pi}_H^R)) \circ \rho_B \circ \mu_B \text{ ((14))} \\
&= \mu_B \circ (\mu_B \otimes (h \circ \Pi_H^R)) \circ \rho_B \circ \mu_B \text{ ((40) of [3])} \\
&= \mu_B \text{ ((8))}.
\end{aligned}$$

Also, if $h : H \rightarrow B$ is a multiplicative morphism of right H -comodules, the following identity holds

$$h \circ \Pi_H^L = q_B \circ h, \quad (16)$$

because

$$\begin{aligned}
&h \circ \Pi_H^L \\
&= \mu_B \circ (h \otimes (h \circ \lambda_H)) \circ \delta_H \text{ (} h \text{ is multiplicative)} \\
&= q_B \circ h \text{ (} h \text{ is a comodule morphism)}.
\end{aligned}$$

If $h : H \rightarrow B$ is an anchor morphism we have that the equality

$$\mu_B \circ ((\mu_B \circ (B \otimes (h \circ \Pi_H^L))) \otimes h) \circ (B \otimes \delta_H) = \mu_B \circ (B \otimes h) \quad (17)$$

holds. Indeed:

$$\begin{aligned}
&\mu_B \circ ((\mu_B \circ (B \otimes (h \circ \Pi_H^L))) \otimes h) \circ (B \otimes \delta_H) \\
&= \mu_B \circ ((\mu_B \circ (\mu_B \otimes (h \circ \lambda_H))) \circ (B \otimes \rho_B)) \otimes h) \circ (B \otimes (\rho_B \circ h)) \text{ ((13) and comodule morphism condition (13) for } h) \\
&= \mu_B \circ ((\mu_B \circ (B \otimes (h \circ \lambda_H))) \otimes h) \circ (B \otimes \delta_H) \circ (\mu_B \otimes H) \circ (B \otimes (\rho_B \circ h)) \text{ (comodule condition for } B) \\
&= \mu_B \circ ((\mu_B \circ (h \circ \Pi_H^R)) \circ (B \otimes (\rho_B \circ h))) \text{ ((c2) of Definition 2.7)} \\
&= \mu_B \circ (B \otimes h) \text{ ((15))}.
\end{aligned}$$

Then, by (16) and the comodule morphism condition for h , (17) is equivalent to

$$\mu_B \circ ((\mu_B \circ (B \otimes q_B)) \otimes h) \circ (B \otimes (\rho_B \circ h)) = \mu_B \circ (B \otimes h). \quad (18)$$

Finally, for an anchor morphism $h : H \rightarrow B$ the equality

$$\mu_B \circ (q_B \otimes h) \circ \rho_B = id_B \quad (19)$$

holds, because

$$\begin{aligned}
&\mu_B \circ (q_B \otimes h) \circ \rho_B \\
&= \mu_B \circ ((\mu_B \circ (B \otimes (h \circ \lambda_H))) \otimes h) \circ (B \otimes \delta_H) \circ \rho_B \text{ (comodule condition for } B) \\
&= \mu_B \circ (B \otimes (h \circ \Pi_H^R)) \circ \rho_B \text{ ((c2) of Definition 2.7)} \\
&= id_B \text{ ((8))}.
\end{aligned}$$

Definition 2.8. Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism and let M be an object in $\mathcal{C}$. We say that (M, ϕ_M, ρ_M) is a strong (H, B, h) -Hopf module if the following axioms hold:

(d1) The pair (M, ρ_M) is a right H -comodule.

(d2) The morphism $\phi_M : M \otimes B \rightarrow M$ satisfies:

$$(d2-1) \quad \phi_M \circ (M \otimes \eta_B) = id_M.$$

$$(d2-2) \quad \phi_M \circ ((\phi_M \circ (M \otimes i_B)) \otimes B) = \phi_M \circ (M \otimes (\mu_B \circ (i_B \otimes B))).$$

$$(d2-3) \quad \rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ (\rho_M \otimes \rho_B).$$

$$(d2-4) \quad \phi_M \circ ((\phi_M \circ (M \otimes h)) \otimes (h \circ \lambda_H)) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes (h \circ \Pi_H^L)).$$

$$(d2-5) \quad \phi_M \circ ((\phi_M \circ (M \otimes (h \circ \lambda_H))) \otimes h) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes (h \circ \Pi_H^R)).$$

For example, the triple (H, μ_H, δ_H) is a strong (H, H, id_H) -Hopf module. Also, if the following equality

$$\mu_B \circ ((\mu_B \circ (B \otimes i_B)) \otimes B) = \mu_B \circ (B \otimes (\mu_B \circ (i_B \otimes B))) \quad (20)$$

holds, the triple (B, μ_B, ρ_B) is a strong (H, B, h) -Hopf module.

Let (M, ϕ_M, ρ_M) be a strong (H, B, h) -Hopf module. In a similar way to (14), but using (d2-3) of Definition 2.8 instead of (4), it is easy to see that

$$(M \otimes \overline{\Pi}_H^R) \circ \rho_M \circ \phi_M = (\phi_M \otimes \overline{\Pi}_H^R) \circ (M \otimes \rho_M) \quad (21)$$

holds. Moreover, by (7), (d2-3), (d2-1), and the condition of comodule for M , we also have the equality

$$\phi_M \circ (M \otimes (h \circ \Pi_H^R)) \circ \rho_M = id_M. \quad (22)$$

As a consequence, we can obtain the Hopf module version of (15), i.e.,

$$\phi_M \circ (\phi_M \otimes (h \circ \Pi_H^R)) \circ (M \otimes \rho_B) = \phi_M. \quad (23)$$

Moreover, in this setting, the equality

$$\phi_M \circ ((\phi_M \circ (M \otimes (h \circ \Pi_H^L))) \otimes h) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes h) \quad (24)$$

holds because

$$\begin{aligned} & \phi_M \circ ((\phi_M \circ (M \otimes (h \circ \Pi_H^L))) \otimes h) \circ (M \otimes \delta_H) \\ &= \phi_M \circ ((\phi_M \circ (M \otimes (q_B \circ h))) \otimes h) \circ (M \otimes \delta_H) \quad ((16)) \\ &= \phi_M \circ (M \otimes (\mu_B \circ ((q_B \circ h) \otimes h) \circ \delta_H)) \quad ((d2-2) \text{ of Definition 2.8}) \\ &= \phi_M \circ (M \otimes (\mu_B \circ ((h \circ \Pi_H^L) \otimes h) \circ \delta_H)) \quad ((16)) \\ &= \phi_M \circ (M \otimes (h \circ (\Pi_H^L * id_H))) \quad (h \text{ is multiplicative}) \\ &= \phi_M \circ (M \otimes h) \quad ((4) \text{ of [3]}). \end{aligned}$$

Finally, by (16) the equality (24) is equivalent to

$$\phi_M \circ ((\phi_M \circ (M \otimes q_B)) \otimes h) \circ (M \otimes (\rho_B \circ h)) = \phi_M \circ (M \otimes h). \quad (25)$$

Proposition 2.9. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism and let (M, ϕ_M, ρ_M) be a strong (H, B, h) -Hopf module. The endomorphism $q_M := \phi_M \circ (M \otimes (h \circ \lambda_H)) \circ \rho_M : M \rightarrow M$ satisfies*

$$\rho_M \circ q_M = (M \otimes \Pi_H^L) \circ \rho_M \circ q_M, \quad (26)$$

$$\rho_M \circ q_M = (M \otimes \overline{\Pi}_H^R) \circ \rho_M \circ q_M, \quad (27)$$

and, as a consequence, q_M is an idempotent. Moreover, if M^{coH} (object of coinvariants) is the image of q_M and $p_M : M \rightarrow M^{coH}$, $i_M : M^{coH} \rightarrow M$ are the morphisms such that $q_M = i_M \circ p_M$ and $id_{M^{coH}} = p_M \circ i_M$,

$$\begin{array}{ccccc} M^{coH} & \xrightarrow{i_M} & M & \xrightarrow{\rho_M} & M \otimes H, \\ & & & \xrightarrow{(M \otimes \overline{\Pi}_H^R) \circ \rho_M} & \\ M^{coH} & \xrightarrow{i_M} & M & \xrightarrow{\rho_M} & M \otimes H, \\ & & & \xrightarrow{(M \otimes \Pi_H^L) \circ \rho_M} & \end{array}$$

are equalizer diagrams.

Proof. The proof is similar to the one developed in Proposition 2.5 but using (d2-3) instead of (4) and the comodule condition for M instead of the comodule condition for B . $\square$

Remark 2.10. Note that, as in the case of B , the object of coinvariants M^{coH} is independent of the anchor morphism h because it is the equalizer object of ρ_M and $(M \otimes \overline{\Pi}_H^R) \circ \rho_M$ (or ρ_M and $(M \otimes \Pi_H^L) \circ \rho_M$).

Lemma 2.11. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism and let (M, ϕ_M, ρ_M) be a strong (H, B, h) -Hopf module. The following equalities hold:*

$$\rho_M \circ \phi_M \circ (i_M \otimes B) = (\phi_M \otimes H) \circ (i_M \otimes \rho_B), \quad (28)$$

$$q_M \circ \phi_M \circ (i_M \otimes B) = \phi_M \circ (i_M \otimes q_B), \quad (29)$$

$$(q_M \otimes H) \circ \rho_M \circ \phi_M \circ (i_M \otimes B) = ((\phi_M \circ (M \otimes q_B)) \otimes H) \circ (i_M \otimes \rho_B), \quad (30)$$

$$(p_M \otimes H) \circ \rho_M \circ \phi_M \circ (i_M \otimes B) = ((p_M \circ \phi_M \circ (M \otimes q_B)) \otimes H) \circ (i_M \otimes \rho_B), \quad (31)$$

$$q_M \circ \phi_M \circ (i_M \otimes i_B) = \phi_M \circ (i_M \otimes i_B), \quad (32)$$

$$p_M \circ \phi_M \circ (i_M \otimes B) = p_M \circ \phi_M \circ (i_M \otimes q_B). \quad (33)$$

$$\phi_M \circ (q_M \otimes h) \circ \rho_M = id_M. \quad (34)$$

Proof. The proof for the first equality is the following:

$$\begin{aligned} & \rho_M \circ \phi_M \circ (i_M \otimes B) \\ &= (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ ((\rho_M \circ i_M) \otimes \rho_B) \text{ ((d2-3) of Definition (2.8))} \\ &= (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ (((M \otimes \bar{\Pi}_H^R) \circ \rho_M \circ i_M) \otimes \rho_B) \text{ ((27))} \\ &= (\phi_M \otimes (\mu_H \circ (\bar{\Pi}_H^R \otimes H))) \circ (M \otimes c_{H,B} \otimes H) \circ ((\rho_M \circ i_M) \otimes \rho_B) \text{ (naturality of } c) \\ &= (\phi_M \otimes (\varepsilon_H \circ \mu_H) \otimes H) \circ (M \otimes c_{H,B} \otimes H \otimes H) \circ ((\rho_M \circ i_M) \otimes ((B \otimes \delta_H) \circ \rho_B)) \text{ ((10) of [3])} \\ &= (((\phi_M \otimes (\varepsilon_H \circ \mu_H)) \circ (M \otimes c_{H,B} \otimes H) \circ (\rho_M \otimes \rho_B)) \otimes H) \circ (i_M \otimes \rho_B) \text{ (comodule condition for } B) \\ &= (((M \otimes \varepsilon_H) \circ \rho_M \circ \phi_M) \otimes H) \circ (i_M \otimes \rho_B) \text{ ((d2-3) of Definition (2.8))} \\ &= (\phi_M \otimes H) \circ (i_M \otimes \rho_B) \text{ (comodule condition for } B). \end{aligned}$$

The equality (29) follows by

$$\begin{aligned} & q_M \circ \phi_M \circ (i_M \otimes B) \\ &= \phi_M \circ (\phi_M \otimes (h \circ \lambda_H)) \circ (i_M \otimes \rho_B) \text{ ((28))} \\ &= \phi_M \circ (i_M \otimes q_B) \text{ ((d2-4) of Definition (2.8))}. \end{aligned}$$

Thus, composing in (28) with $q_M \otimes H$, we obtain (30). Also, composing in (30) with $p_M \otimes B$, we have (31). Moreover, composing with $M^{coH} \otimes i_B$ in (29), we obtain (32), and doing the same with p_M we get (33). Finally, (34) holds because

$$\begin{aligned} & \phi_M \circ (q_M \otimes h) \circ \rho_M \\ &= \phi_M \circ ((\phi_M \circ (M \otimes (h \circ \lambda_H))) \otimes h) \circ (M \otimes \delta_H) \circ \rho_M \text{ (comodule condition for } M) \\ &= \phi_M \circ (M \otimes (h \circ \Pi_H^R)) \circ \rho_M \text{ ((d2-5) of Definition (2.8))} \\ &= id_M \text{ ((22))}. \end{aligned}$$

□

Proposition 2.12. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism. If (20) holds, the submagma of coinvariants $(B^{coH}, \eta_{B^{coH}}, \mu_{B^{coH}})$ is a monoid.*

Proof. Firstly, remember that if (20) holds, the triple (B, μ_B, ρ_B) is a strong (H, B, h) -Hopf module. Then by (29),

$$q_B \circ \mu_B \circ (i_B \otimes i_B) = \mu_B \circ (i_B \otimes i_B). \quad (35)$$

As a consequence, B^{coH} is a monoid because,

$$\begin{aligned} & i_B \circ \mu_{B^{coH}} \circ (B^{coH} \otimes \mu_{B^{coH}}) \\ &= q_B \circ \mu_B \circ (i_B \otimes (q_B \circ \mu_B \circ (i_B \otimes i_B))) \text{ ((12))} \\ &= q_B \circ \mu_B \circ (i_B \otimes (\mu_B \circ (i_B \otimes i_B))) \text{ ((35))} \\ &= q_B \circ \mu_B \circ ((\mu_B \circ (i_B \otimes i_B)) \otimes i_B) \text{ ((20))} \\ &= q_B \circ \mu_B \circ ((q_B \circ \mu_B \circ (i_B \otimes i_B)) \otimes i_B) \text{ ((35))} \\ &= i_B \circ \mu_{B^{coH}} \circ (\mu_{B^{coH}} \otimes B^{coH}) \text{ ((12))}. \end{aligned}$$

□

Proposition 2.13. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism. If (20) holds, for all strong (H, B, h) -Hopf module (M, ϕ_M, ρ_M) , the object of coinvariants M^{coH} is a right B^{coH} -module.*

Proof. First note that, by Proposition 2.12, the object B^{coH} is a monoid. Now we will show that there exists an action $\phi_{M^{coH}} : M^{coH} \otimes B^{coH} \rightarrow M^{coH}$ such that $\phi_{M^{coH}} \circ (M^{coH} \otimes \eta_{B^{coH}}) = id_{M^{coH}}$, and $\phi_{M^{coH}} \circ (\phi_{M^{coH}} \otimes B^{coH}) = \phi_{M^{coH}} \circ (M^{coH} \otimes \mu_{B^{coH}})$. To define the action, we begin by proving that

$$\rho_M \circ \phi_M \circ (i_M \otimes i_B) = (M \otimes \Pi_H^L) \circ \rho_M \circ \phi_M \circ (i_M \otimes i_B). \quad (36)$$

Indeed,

$$\begin{aligned} & \rho_M \circ \phi_M \circ (i_M \otimes i_B) \\ &= (\phi_M \otimes H) \circ (i_M \otimes (\rho_B \circ i_B)) \quad ((28)) \\ &= (\phi_M \otimes \Pi_H^L) \circ (i_M \otimes (\rho_B \circ i_B)) \quad ((5)) \\ &= (M \otimes \Pi_H^L) \circ \rho_M \circ \phi_M \circ (i_M \otimes i_B) \quad ((28)) \end{aligned}$$

Then, there exists a unique morphism $\phi_{M^{coH}} : M^{coH} \otimes B^{coH} \rightarrow M^{coH}$ such that

$$\phi_M \circ (i_M \otimes i_B) = i_M \circ \phi_{M^{coH}}. \quad (37)$$

Therefore,

$$\phi_{M^{coH}} = p_M \circ \phi_M \circ (i_M \otimes i_B). \quad (38)$$

The pair $(M^{coH}, \phi_{M^{coH}})$ satisfies the conditions of right B^{coH} -module because

$$\phi_{M^{coH}} \circ (M^{coH} \otimes \eta_{B^{coH}}) \stackrel{(9)}{=} p_M \circ \phi_M \circ (i_M \otimes \eta_B) \stackrel{(d2-1)}{=} p_M \circ i_M = id_{M^{coH}},$$

and

$$\begin{aligned} & \phi_{M^{coH}} \circ (\phi_{M^{coH}} \otimes B^{coH}) \\ &= p_M \circ \phi_M \circ ((q_M \circ \phi_M \circ (i_M \otimes i_B)) \otimes i_B) \quad ((38)) \\ &= p_M \circ \phi_M \circ ((\phi_M \circ (i_M \otimes i_B)) \otimes i_B) \quad ((32)) \\ &= p_M \circ \phi_M \circ (i_M \otimes (\mu_B \circ (i_B \otimes i_B))) \quad ((d2-2) \text{ of Definition (2.8)}) \\ &= p_M \circ \phi_M \circ (i_M \otimes (i_B \circ \mu_{B^{coH}})) \quad ((10)) \\ &= \phi_{M^{coH}} \circ (M^{coH} \otimes \mu_{B^{coH}}) \quad ((38)). \end{aligned}$$

□

Proposition 2.14. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism. Assume that (20) and*

$$\mu_B \circ (i_B \otimes \mu_B) = \mu_B \circ ((\mu_B \circ (i_B \otimes B)) \otimes B) \quad (39)$$

hold. Then if the category $\mathcal{C}$ admits coequalizers and the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers, for all strong (H, B, h) -Hopf module (M, ϕ_M, ρ_M) , the object $M^{coH} \otimes_{B^{coH}} B$, defined by the coequalizer of $T_M^1 = \phi_{M^{coH}} \otimes B$ and $T_M^2 = M^{coH} \otimes (\mu_B \circ (i_B \otimes B))$, is a strong (H, B, h) -Hopf module. Moreover, $M^{coH} \otimes_{B^{coH}} B$ and M are isomorphic as right H -comodules.

Proof. In the first step, we begin by proving the existence of an action and a coaction for $M^{coH} \otimes_{B^{coH}} H$. If the object $M^{coH} \otimes_{B^{coH}} H$ is defined by the coequalizer diagram

$$\begin{array}{ccc} M^{coH} \otimes B^{coH} \otimes B & \xrightarrow{T_M^1} & M^{coH} \otimes B \xrightarrow{n_{M^{coH}}} M^{coH} \otimes_{B^{coH}} B, \\ & \xrightarrow{T_M^2} & \end{array} \quad (40)$$

we have that

$$\begin{array}{ccc} M^{coH} \otimes B^{coH} \otimes B \otimes B & \xrightarrow{T_M^1 \otimes B} & M^{coH} \otimes B \otimes B \xrightarrow{n_{M^{coH}} \otimes B} M^{coH} \otimes_{B^{coH}} B \otimes B \\ & \xrightarrow{T_M^2 \otimes B} & \end{array} \quad (41)$$

is also a coequalizer diagram because the functor $- \otimes B$ preserves coequalizers. Consider the morphism $n_{M^{coH}} \circ (M^{coH} \otimes \mu_B) : M^{coH} \otimes B \otimes B \rightarrow M^{coH} \otimes_{B^{coH}} B$. Then,

$$n_{M^{coH}} \circ (\phi_{M^{coH}} \otimes \mu_B) \stackrel{(40)}{=} n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (i_B \otimes \mu_B))) \stackrel{(39)}{=} n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ ((\mu_B \circ (i_B \otimes B)) \otimes B))),$$

and, as a consequence, there exists a unique morphism

$$\phi_{M^{coH} \otimes_{B^{coH}} B} : M^{coH} \otimes_{B^{coH}} B \otimes B \rightarrow M^{coH} \otimes_{B^{coH}} B$$

such that

$$\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes B) = n_{M^{coH}} \circ (M^{coH} \otimes \mu_B). \quad (42)$$

On the other hand, consider the morphism

$$(n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes \rho_B) : M^{coH} \otimes B \rightarrow M^{coH} \otimes_{B^{coH}} B \otimes H.$$

Then, taking into account that, by (20), (B, μ_B, ρ_B) is a strong (H, B, h) -Hopf module,

$$\begin{aligned} & (n_{M^{coH}} \otimes H) \circ (\phi_{M^{coH}} \otimes \rho_B) \\ &= (n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes ((\mu_B \otimes H) \circ (i_B \otimes \rho_B))) \stackrel{(40)}{=} \\ &= (n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes (\rho_B \circ \mu_B \circ (i_B \otimes B))) \stackrel{(28)}{=} \end{aligned}$$

Thus, there exists a unique morphism

$$\rho_{M^{coH} \otimes_{B^{coH}} B} : M^{coH} \otimes_{B^{coH}} B \rightarrow M^{coH} \otimes_{B^{coH}} B \otimes H$$

such that

$$\rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}} = (n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes \rho_B). \quad (43)$$

We proceed to show that $(M^{coH} \otimes_{B^{coH}} B, \phi_{M^{coH} \otimes_{B^{coH}} B}, \rho_{M^{coH} \otimes_{B^{coH}} B})$ is a strong (H, B, h) -Hopf module. Indeed, first note that by (43) and the comodule condition for B ,

$$(M^{coH} \otimes_{B^{coH}} B \otimes \varepsilon_H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}} = (n_{M^{coH}} \otimes \varepsilon_H) \circ (M^{coH} \otimes \rho_B) = n_{M^{coH}}.$$

As a consequence, and using that $n_{M^{coH}}$ is a coequalizer, we obtain that

$$(M^{coH} \otimes_{B^{coH}} B \otimes \varepsilon_H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} = id_{M^{coH} \otimes_{B^{coH}} B}.$$

By similar arguments we have

$$\begin{aligned} & (\rho_{M^{coH} \otimes_{B^{coH}} B} \otimes H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}} = (n_{M^{coH}} \otimes H \otimes H) \circ (M^{coH} \otimes ((\rho_B \otimes H) \circ \rho_B)) \\ &= (n_{M^{coH}} \otimes H \otimes H) \circ (M^{coH} \otimes ((B \otimes \delta_H) \circ \rho_B)) = (M^{coH} \otimes_{B^{coH}} B \otimes \delta_H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}}. \end{aligned}$$

Therefore, the pair $(M^{coH} \otimes_{B^{coH}} B, \rho_{M^{coH} \otimes_{B^{coH}} B})$ is a right H -comodule and we have (d1) of Definition 2.8.

On the other hand, by (42),

$$\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes \eta_B) = n_{M^{coH}},$$

and then, (d2-1) of Definition 2.8, i.e., $\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (M^{coH} \otimes_{B^{coH}} B \otimes \eta_B) = id_{M^{coH} \otimes_{B^{coH}} B}$, holds. Moreover,

$$\begin{aligned} & \phi_{M^{coH} \otimes_{B^{coH}} B} \circ ((\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes q_B)) \otimes B) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ ((\mu_B \circ (B \otimes q_B)) \otimes B))) \stackrel{(42)}{=} \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (B \otimes (\mu_B \circ (q_B \otimes B)))) \stackrel{(20)}{=} \\ &= \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes (\mu_B \circ (q_B \otimes B))) \stackrel{(42)}{=} \end{aligned}$$

and, as a consequence,

$$\begin{aligned} & \phi_{M^{coH} \otimes_{B^{coH}} B} \circ ((\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (M^{coH} \otimes_{B^{coH}} B \otimes q_B)) \otimes B) \\ &= \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (M^{coH} \otimes_{B^{coH}} B \otimes (\mu_B \circ (q_B \otimes B))) \end{aligned}$$

because $n_{M^{coH}} \otimes B \otimes B$ is a coequalizer. Therefore (d2-2) of Definition 2.8 follows because p_B is a projection.

The proof for (d2-3) of Definition 2.8 is the following: Composing with the coequalizer $n_{M^{coH}} \otimes B$ we obtain that

$$\begin{aligned} & \rho_{M^{coH} \otimes_{B^{coH}} B} \circ \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes B) \\ &= (n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes (\rho_B \circ \mu_B)) \quad ((42), (43)) \\ &= (n_{M^{coH}} \otimes H) \circ (M^{coH} \otimes (\mu_B \otimes H \circ (\rho_B \otimes \rho_B))) \quad ((4)) \\ &= (\phi_{M^{coH} \otimes_{B^{coH}} B} \otimes \mu_H) \circ (M^{coH} \otimes_{B^{coH}} B \otimes c_{H,B} \otimes H) \circ ((\rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}}) \otimes \rho_B) \quad ((42), (43)) \end{aligned}$$

and thus

$$\rho_{M^{coH} \otimes_{B^{coH}} B} \circ \phi_{M^{coH} \otimes_{B^{coH}} B} = (\phi_{M^{coH} \otimes_{B^{coH}} B} \otimes \mu_H) \circ (M^{coH} \otimes_{B^{coH}} B \otimes c_{H,B} \otimes H) \circ (\rho_{M^{coH} \otimes_{B^{coH}} B} \otimes \rho_B).$$

Using that $- \otimes H$ preserves coequalizers, the equality (d2-4) of Definition 2.8 follows from

$$\begin{aligned} & \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (\phi_{M^{coH} \otimes_{B^{coH}} B} \otimes B) \circ (n_{M^{coH}} \otimes ((h \otimes (h \circ \lambda_H)) \circ \delta_H)) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (\mu_B \otimes B) \circ (B \otimes ((h \otimes (h \circ \lambda_H)) \circ \delta_H)))) \quad ((42)) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (B \otimes (h \circ \Pi_H^L)))) \quad ((c1) \text{ of Definition 2.7}) \\ &= \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes (h \circ \Pi_H^L)) \quad ((42)). \end{aligned}$$

Also, we have

$$\begin{aligned} & \phi_{M^{coH} \otimes_{B^{coH}} B} \circ ((\phi_{M^{coH} \otimes_{B^{coH}} B} \circ (M^{coH} \otimes_{B^{coH}} B \otimes (h \circ \lambda_H))) \otimes h) \circ (n_{M^{coH}} \otimes \delta_H) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ ((\mu_B \circ (B \otimes (h \circ \lambda_H))) \otimes h) \circ (B \otimes \delta_H))) \quad ((42)) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (B \otimes (h \circ \Pi_H^R)))) \quad ((c2) \text{ of Definition 2.7}) \\ &= \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes (h \circ \Pi_H^R)) \quad ((42)), \end{aligned}$$

and then (d2-5) of Definition 2.8 holds.

Consider the morphism $\phi_M \circ (i_M \otimes B) : M^{coH} \otimes B \rightarrow M$. By (37) and (d2-2) of Definition 2.8, we obtain that $\phi_M \circ (i_M \otimes B) \circ T_M^1 = \phi_M \circ (i_M \otimes B) \circ T_M^2$ and, as a consequence, there exists a unique morphism $\omega_M : M^{coH} \otimes_{B^{coH}} B \rightarrow M$ such that

$$\omega_M \circ n_{M^{coH}} = \phi_M \circ (i_M \otimes B). \quad (44)$$

The morphism ω_M is a morphism of right H -comodules because

$$\begin{aligned} (\omega_M \otimes H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} \circ n_{M^{coH}} &\stackrel{(43)}{=} ((\omega_M \circ n_{M^{coH}}) \otimes H) \circ (M^{coH} \otimes \rho_B) \stackrel{(44)}{=} (\phi_M \otimes H) \circ (i_M \otimes \rho_B) \\ &\stackrel{(28)}{=} \rho_M \circ \phi_M \circ (i_M \otimes B) \stackrel{(44)}{=} \rho_M \circ \omega_M \circ n_{M^{coH}} \end{aligned}$$

and then $(\omega_M \otimes H) \circ \rho_{M^{coH} \otimes_{B^{coH}} B} = \rho_M \circ \omega_M$.

Finally, ω_M is an isomorphism with inverse $\omega'_M = n_{M^{coH}} \circ (p_M \otimes h) \circ \rho_M$. Indeed, firstly note that, by (34), we have that $\omega_M \circ \omega'_M = id_M$. On the other hand, composing with $n_{M^{coH}}$ we have

$$\begin{aligned} & \omega'_M \circ \omega_M \circ n_{M^{coH}} \\ &= n_{M^{coH}} \circ (p_M \otimes h) \circ \rho_M \circ \phi_M \circ (i_M \otimes B) \quad ((44)) \\ &= n_{M^{coH}} \circ ((p_M \circ \phi_M \circ (M \otimes q_B)) \otimes h) \circ (i_M \otimes \rho_B) \quad ((31)) \\ &= n_{M^{coH}} \circ ((\phi_{M^{coH}} \circ (M \otimes p_B)) \otimes h) \circ (M^{coH} \otimes \rho_B) \quad ((38)) \\ &= n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (q_B \otimes h) \circ \rho_B)) \quad ((40)) \\ &= n_{M^{coH}} \quad ((19)), \end{aligned}$$

and, as a consequence, $\omega'_M \circ \omega_M = id_{M^{coH} \otimes_{B^{coH}} B}$. □

Remark 2.15. Note that in the previous proposition the existence of the comodule structure on $M^{coH} \otimes_{B^{coH}} B$ does not depend on the preservation of coequalizers by the functors $- \otimes B$ and $- \otimes H$.

Proposition 2.16. *Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism and let (P, ϕ_P, ρ_P) , (Q, ϕ_Q, ρ_Q) be strong (H, B, h) -Hopf modules. If there exists a right H -comodule isomorphism $\omega : Q \rightarrow P$, the triple $(P, \phi_P^\omega = \omega \circ \phi_Q \circ (\omega^{-1} \otimes B), \rho_P)$, called the ω -deformation of (P, ϕ_P, ρ_P) , is a strong (H, B, h) -Hopf module.*

Proof. The proof follows easily because, if ω is a right H -comodule isomorphism, $\rho_P = (\omega \otimes H) \circ \rho_Q \circ \omega^{-1}$. $\square$

Definition 2.17. Let H be a weak Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma. Let $h : H \rightarrow B$ be an anchor morphism. Assume that (20) and (39) hold. Then if the category $\mathcal{C}$ admits coequalizers and the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers, we define the category of strong (H, B, h) -Hopf modules as the one whose objects are strong (H, B, h) -Hopf modules, and whose morphisms $f : M \rightarrow N$ are morphisms of right H -comodules and B -quasilinear, i.e.

$$\phi_N^{\omega_N} \circ (f \otimes B) = f \circ \phi_M^{\omega_M}, \quad (45)$$

where $\omega_M : M^{coH} \otimes_{B^{coH}} B \rightarrow M$, $\omega_N : N^{coH} \otimes_{B^{coH}} B \rightarrow N$ are the isomorphisms of right H -comodules obtained in the proof of Proposition 2.14. This category will be denoted by $\mathcal{SM}_B^H(h)$.

Example 2.18. As particular instances of $\mathcal{SM}_B^H(h)$ we will describe in detail some interesting examples of Hopf module categories associated to Hopf algebras, weak Hopf algebras, Hopf quasigroups and weak Hopf quasigroups.

Examples for Hopf algebras: If H is a Hopf algebra and $B = H$, $\rho_B = \delta_H$, $h = id_H$, the equalities (20) and (39) hold trivially because the product on H is associative. In this case the category $\mathcal{SM}_H^H(id_H)$ is the one whose objects are triples (M, ϕ_M, ρ_M) where:

- (e1) The pair (M, ρ_M) is a right H -comodule.
- (e2) The morphism $\phi_M : M \otimes H \rightarrow M$ satisfies:
 - (e2-1) $\phi_M \circ (M \otimes \eta_H) = id_M$,
 - (e2-2) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,H} \otimes H) \circ (\rho_M \otimes \delta_H)$,
 - (e2-3) $\phi_M \circ (\phi_M \otimes \lambda_H) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,
 - (e2-4) $\phi_M \circ ((\phi_M \circ (M \otimes \lambda_H)) \otimes H) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,

because in this case $\Pi_H^L = \Pi_H^R = q_H = \eta_H \otimes \varepsilon_H$, $H^{coH} = K$, and $i_H = \eta_H$. Then, if (M, ϕ_M, ρ_M) is a classical Hopf module in the sense of Larson and Sweedler [21] (see also [30]), we have that (M, ϕ_M, ρ_M) is an object in $\mathcal{SM}_H^H(id_H)$ because the identity

$$\phi_M \circ (\phi_M \otimes H) = \phi_M \circ (M \otimes \mu_H) \quad (46)$$

holds. Moreover, the morphisms of $\mathcal{SM}_H^H(id_H)$ are morphisms of right H -comodules and H -quasilinear, i.e., satisfying (45), where $\omega_M = \phi_M \circ (i_M \otimes H) : M^{coH} \otimes H \rightarrow M$ is the associated isomorphism with inverse $\omega_M^{-1} = (p_M \otimes H) \circ \rho_M$, and $\phi_{M^{coH} \otimes H} = M^{coH} \otimes \mu_H$ and $\rho_{M^{coH} \otimes H} = M^{coH} \otimes \delta_H$. Then any morphism $f : M \rightarrow N$ of right H -comodules and linear in the classical sense, i.e., such that $f \circ \phi_M = \phi_N \circ (f \otimes H)$, satisfies (45). Therefore, the Larson-Sweedler category of Hopf modules, denoted by $\mathcal{M}_H^H$, is a subcategory of $\mathcal{SM}_H^H(id_H)$.

Also, in the Hopf algebra setting we have the following more general example: Let (B, ρ_B) be a right H -comodule monoid such that the functors $- \otimes B$, $- \otimes H$ preserve coequalizers. In this case, the equalities (20) and (39) hold trivially because the product on B is associative. For any h multiplicative total integral, therefore an anchor morphism, the category $\mathcal{SM}_B^H(h)$ is the one whose objects are triples (M, ϕ_M, ρ_M) where:

- (f1) The pair (M, ρ_M) is a right H -comodule.
- (f2) The morphism $\phi_M : M \otimes B \rightarrow M$ satisfies:
 - (f2-1) $\phi_M \circ (M \otimes \eta_B) = id_M$,
 - (f2-2) $\phi_M \circ ((\phi_M \circ (M \otimes i_B)) \otimes B) = \phi_M \circ (M \otimes (\mu_B \circ (i_B \otimes B)))$,
 - (f2-3) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ (\rho_M \otimes \rho_B)$,
 - (f2-4) $\phi_M \circ ((\phi_M \circ (M \otimes h)) \otimes (h \circ \lambda_H)) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,
 - (f2-5) $\phi_M \circ ((\phi_M \circ (M \otimes (h \circ \lambda_H))) \otimes h) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,

because in this case $\Pi_H^L = \Pi_H^R = \eta_H \otimes \varepsilon_H$. The morphisms in $\mathcal{SM}_B^H(h)$ are morphisms of right H -comodules and B -quasilinear where $\omega_M : M^{coH} \otimes_{B^{coH}} H \rightarrow M$ is the associated isomorphism of right

H -comodules defined in the proof of Proposition 2.14. Then, if (M, ϕ_M, ρ_M) is an (H, B) -Hopf module in the sense of Doi [13], we have that (M, ϕ_M, ρ_M) is an object in $\mathcal{SM}_B^H(h)$ because the identity

$$\phi_M \circ (\phi_M \otimes B) = \phi_M \circ (M \otimes \mu_B) \quad (47)$$

holds. The morphisms in $\mathcal{SM}_B^H(h)$ are colinear morphisms satisfying (45), for the action $\phi_{M \circ H \otimes_{B^{coH}} H}$ and the coaction $\rho_{M \circ H \otimes_{B^{coH}} H}$ defined in the proof of Proposition 2.14. Then any morphism $f : M \rightarrow N$ of right H -comodules and right B -linear, i.e., such that $f \circ \phi_M = \phi_N \circ (f \otimes B)$, satisfies (45). Therefore, the category of right (H, B) -Hopf modules, denoted by $\mathcal{M}_B^H$, is a subcategory of $\mathcal{SM}_B^H(h)$ (for all multiplicative total integral h).

Examples for weak Hopf algebras: Let H be a weak Hopf algebra. Let (B, ρ_B) be a right H -comodule monoid such that the functors $- \otimes B$, $- \otimes H$ preserve coequalizers. As in the Hopf algebra setting, the equalities (20) and (39) hold trivially because the product on B is associative. Also, in this case, a multiplicative total integral h is an anchor morphism because the product on B is associative, (7) holds and $\Pi_H^L * id_H = id_H$. Then, the category $\mathcal{SM}_B^H(h)$ is the one whose objects are triples (M, ϕ_M, ρ_M) where:

- (g1) The pair (M, ρ_M) is a right H -comodule.
- (g2) The morphism $\phi_M : M \otimes B \rightarrow M$ satisfies:
 - (g2-1) $\phi_M \circ (M \otimes \eta_B) = id_M$,
 - (g2-2) $\phi_M \circ ((\phi_M \circ (M \otimes i_B)) \otimes B) = \phi_M \circ (M \otimes (\mu_B \circ (i_B \otimes B)))$,
 - (g2-3) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ (\rho_M \otimes \rho_B)$,
 - (g2-4) $\phi_M \circ ((\phi_M \circ (M \otimes h)) \otimes (h \circ \lambda_H)) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes (h \circ \Pi_H^L))$,
 - (g2-5) $\phi_M \circ ((\phi_M \circ (M \otimes (h \circ \lambda_H))) \otimes h) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes (h \circ \Pi_H^R))$.

Then, if (M, ϕ_M, ρ_M) is an (H, B) -Hopf module in the sense Böhm [9] (see also [33]), the triple (M, ϕ_M, ρ_M) is an object in $\mathcal{SM}_B^H(h)$ because the identity (47) holds.

As in the Hopf case, the morphisms in $\mathcal{SM}_B^H(h)$ are colinear morphisms satisfying (45), for $\phi_{M \circ H \otimes_{B^{coH}} H}$ and $\rho_{M \circ H \otimes_{B^{coH}} H}$ the action and the coaction defined in the proof of Proposition 2.14. Then any morphism $f : M \rightarrow N$ of right H -comodules and right B -linear satisfies (45). Therefore, as in the classical context, the category of right (H, B) -Hopf modules, denoted by $\mathcal{M}_B^H$, is a subcategory of $\mathcal{SM}_B^H(h)$ (for all multiplicative total integral h). As a consequence, if $H = B$, $\rho_B = \delta_H$ and $h = id_H$, we obtain that the category $\mathcal{M}_H^H$, i.e., the category of Hopf modules associated to H , is a subcategory of $\mathcal{SM}_H^H(id_H)$. Note that in this case the objects of $\mathcal{SM}_H^H(id_H)$ are triples (M, ϕ_M, ρ_M) where:

- (h1) The pair (M, ρ_M) is a right H -comodule.
- (h2) The morphism $\phi_M : M \otimes H \rightarrow M$ satisfies:
 - (h2-1) $\phi_M \circ (M \otimes \eta_H) = id_M$,
 - (h2-2) $\phi_M \circ ((\phi_M \circ (M \otimes i_L)) \otimes H) = \phi_M \circ (M \otimes (\mu_H \circ (i_L \otimes H)))$,
 - (h2-3) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,H} \otimes H) \circ (\rho_M \otimes \delta_H)$,
 - (h2-4) $\phi_M \circ (\phi_M \otimes \lambda_H) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes \Pi_H^L)$.
 - (h2-5) $\phi_M \circ ((\phi_M \circ (M \otimes \lambda_H)) \otimes H) \circ (M \otimes \delta_H) = \phi_M \circ (M \otimes \Pi_H^R)$,

because in this case $B^{coH} = H_L$ and $i_B = i_L$.

Examples for Hopf quasigroups: The following example comes from the nonassociative setting. Let H be a Hopf quasigroup and assume that $B = H$, $\rho_B = \delta_H$, $h = id_H$. In this case the equalities (20) and (39) hold because $q_H = \eta_H \otimes \varepsilon_H$, $H^{coH} = K$, and $i_H = \eta_H$. Then the category $\mathcal{SM}_H^H(id_H)$ is the one whose objects are triples (M, ϕ_M, ρ_M) where:

- (i1) The pair (M, ρ_M) is a right H -comodule.
- (i2) The morphism $\phi_M : M \otimes H \rightarrow M$ satisfies:
 - (i2-1) $\phi_M \circ (M \otimes \eta_H) = id_M$,
 - (i2-2) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,H} \otimes H) \circ (\rho_M \otimes \delta_H)$,
 - (i2-3) $\phi_M \circ (\phi_M \otimes (M \circ \lambda_H)) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,
 - (i2-4) $\phi_M \circ (\phi_M \otimes \lambda_H) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$.

Note that in this setting the equality $\phi_M \circ ((\phi_M \circ (M \otimes i_H)) \otimes H) = \phi_M \circ (M \otimes (\mu_H \circ (i_H \otimes H)))$ holds because $i_H = \eta_H$. The morphisms of $\mathcal{SM}_H^H(id_H)$ are morphisms of right H -comodules and H -quasilinear, i.e., satisfying (45), where $\omega_M = \phi_M \circ (i_M \otimes H) : M^{coH} \otimes H \rightarrow M$ is the associated isomorphism with inverse $\omega_M^{-1} = (p_M \otimes H) \circ \rho_M$, and $\phi_{M^{coH} \otimes H} = M^{coH} \otimes \mu_H$ and $\rho_{M^{coH} \otimes H} = M^{coH} \otimes \delta_H$. Therefore $\mathcal{SM}_H^H(id_H)$ is the category of Hopf modules introduced by Brzeziński in [10].

In the previous nonassociative setting, let (B, ρ_B) be a right H -comodule magma such that the functors $-\otimes B$ and $-\otimes H$ preserve coequalizers, and such that the equalities (20) and (39) hold. For any anchor morphism h , the category $\mathcal{SM}_B^H(h)$ is the one whose objects are triples (M, ϕ_M, ρ_M) where:

- (j1) The pair (M, ρ_M) is a right H -comodule.
- (j2) The morphism $\phi_M : M \otimes B \rightarrow M$ satisfies:
 - (j2-1) $\phi_M \circ (M \otimes \eta_B) = id_M$,
 - (j2-2) $\phi_M \circ ((\phi_M \circ (M \otimes i_B)) \otimes B) = \phi_M \circ (M \otimes (\mu_B \circ (i_B \otimes B)))$,
 - (j2-3) $\rho_M \circ \phi_M = (\phi_M \otimes \mu_H) \circ (M \otimes c_{H,B} \otimes H) \circ (\rho_M \otimes \rho_B)$,
 - (j2-4) $\phi_M \circ ((\phi_M \circ (M \otimes h)) \otimes (h \circ \lambda_H)) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,
 - (j2-5) $\phi_M \circ ((\phi_M \circ (M \otimes (h \circ \lambda_H))) \otimes h) \circ (M \otimes \delta_H) = M \otimes \varepsilon_H$,

because in this case $\Pi_H^L = \Pi_H^R = \eta_H \otimes \varepsilon_H$. The morphisms of $\mathcal{SM}_B^H(h)$ are morphisms of right H -comodules and B -quasilinear where $\omega_M : M^{coH} \otimes_{B^{coH}} H \rightarrow M$ is the associated isomorphism of right H -comodules defined in the proof of Proposition 2.14.

Examples for weak Hopf quasigroups: If H is a weak Hopf quasigroup $\mathcal{SM}_H^H(id_H)$ is the category of strong Hopf modules defined in [6] and denoted by $\mathcal{SM}_H^H$. Note that, as a consequence of (24), we obtain that

$$\phi_M \circ (\phi_M \otimes H) \circ (M \otimes \Pi_H^L \otimes H) \circ (M \otimes \delta_H) = \phi_M$$

is a superfluous identity in the definition of strong Hopf module introduced in [6].

Proposition 2.19. *Assume that the conditions of Proposition 2.14 hold. Let (M, ϕ_M, ρ_M) be an object in $\mathcal{SM}_B^H(h)$. Let ω_M be the isomorphism of right H -comodules between $M^{coH} \otimes_{B^{coH}} B$ and M . Then the triple $(M, \phi_M^{\omega_M}, \rho_M)$ is a strong (H, B, h) -Hopf module with the same object of coinvariants that (M, ϕ_M, ρ_M) . Also, the identity*

$$\phi_M^{\omega_M} = \phi_M \circ (q_M \otimes (\mu_B \circ (h \otimes B))) \circ (\rho_M \otimes B) \quad (48)$$

holds and

$$q_M^{\omega_M} = q_M, \quad (49)$$

where $q_M^{\omega_M} = \phi_M^{\omega_M} \circ (M \otimes (h \circ \lambda_H)) \circ \rho_M$ is the idempotent morphism associated to the Hopf module $(M, \phi_M^{\omega_M}, \rho_M)$. Moreover, for $(M, \phi_M^{\omega_M}, \rho_M)$, the associated isomorphism of right H -comodules between $M^{coH} \otimes_{B^{coH}} B$ and M is ω_M , and the equality

$$(\phi_M^{\omega_M})^{\omega_M} = \phi_M^{\omega_M} \quad (50)$$

holds. Finally, there exists an idempotent functor

$$D : \mathcal{SM}_B^H(h) \rightarrow \mathcal{SM}_B^H(h),$$

called the ω -deformation functor, defined on objects by $D((M, \phi_M, \rho_M)) = (M, \phi_M^{\omega_M}, \rho_M)$ and on morphisms by the identity.

Proof. By Proposition 2.16, $(M, \phi_M^{\omega_M}, \rho_M)$ is a strong (H, B, h) -Hopf module. Moreover, note that (48) follows by (42) and (44). Then $q_M^{\omega_M} = q_M$ because

$$\begin{aligned} & q_M^{\omega_M} \\ &= \phi_M \circ (q_M \otimes (\mu_B \circ (h \otimes h))) \circ (\rho_M \otimes \lambda_H) \circ \rho_M \quad ((48)) \\ &= \phi_M \circ (q_M \otimes (h \circ \Pi_H^L)) \circ \rho_M \quad (\text{comodule condition for } M \text{ and multiplicative condition for } h) \\ &= \phi_M \circ (\phi_M \otimes B) \circ (q_M \otimes h \otimes (h \circ \lambda_H)) \circ (M \otimes \delta_H) \circ \rho_M \quad ((d2-4) \text{ of Definition 2.8}) \\ &= \phi_M \circ ((\phi_M \circ (q_M \otimes h)) \circ \rho_M) \otimes (h \circ \lambda_H) \circ \rho_M \quad (\text{comodule condition for } M) \\ &= q_M \quad ((34)). \end{aligned}$$

Therefore, $(M, \phi_M^{\omega_M}, \rho_M)$ has the same object of coinvariants that (M, ϕ_M, ρ_M) . On the other hand,

$$\begin{aligned}
& (\phi_M^{\omega_M})^{\omega_M} \\
&= \phi_M \circ (q_M \otimes \mu_B) \circ (((M \otimes h) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B))) \circ (\rho_M \otimes B) \quad ((48)) \\
&= \phi_M \circ (q_M \otimes \mu_B) \circ (((M \otimes (h \circ \Pi_H^L)) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B))) \circ (\rho_M \otimes B) \quad ((26)) \\
&= \phi_M \circ (q_M \otimes \mu_B) \circ (((M \otimes (q_B \circ h)) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B))) \circ (\rho_M \otimes B) \quad ((16)) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes (q_B \circ h))) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B)) \circ (\rho_M \otimes B) \quad ((d2-2) \text{ of Definition 2.8}) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes (h \circ \Pi_H^L))) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B)) \circ (\rho_M \otimes B) \quad ((16)) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes h)) \circ \rho_M \circ q_M) \otimes (\mu_B \circ (h \otimes B)) \circ (\rho_M \otimes B) \quad ((26)) \\
&= \phi_M^{\omega_M} \quad ((34)).
\end{aligned}$$

Finally, by (49) and (50), it is easy to show that D is a well defined idempotent endofunctor. $\square$

Lemma 2.20. *Assume that the conditions of Proposition 2.14 hold. Let (M, ϕ_M, ρ_M) be an object in $\mathcal{SM}_B^H(h)$. The following identity holds:*

$$\phi_M^{\omega_M} \circ (i_M \otimes B) = \phi_M \circ (i_M \otimes B). \quad (51)$$

Proof. Indeed,

$$\begin{aligned}
& \phi_M^{\omega_M} \circ (i_M \otimes B) \\
&= \phi_M \circ (q_M \otimes \mu_B) \circ (((M \otimes (h \circ \Pi_H^L)) \circ \rho_M \circ i_M) \otimes B) \quad ((26)) \\
&= \phi_M \circ (q_M \otimes \mu_B) \circ (((M \otimes (q_B \circ h)) \circ \rho_M \circ i_M) \otimes B) \quad ((16)) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes (q_B \circ h))) \circ \rho_M \circ i_M) \otimes B \quad ((d2-2) \text{ of Definition 2.8}) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes (h \circ \Pi_H^L))) \circ \rho_M \circ i_M) \otimes B \quad ((16)) \\
&= \phi_M \circ ((\phi_M \circ (q_M \otimes h)) \circ \rho_M \circ i_M) \otimes B \quad ((26)) \\
&= \phi_M \circ (i_M \otimes B) \quad ((34)).
\end{aligned}$$

$\square$

Proposition 2.21. *Assume that the conditions of Proposition 2.14 hold. For any object (M, ϕ_M, ρ_M) in $\mathcal{SM}_B^H(h)$, the strong (H, B, h) -Hopf module $(M^{coH} \otimes_{B^{coH}} B, \phi_{M^{coH} \otimes_{B^{coH}} B}, \rho_{M^{coH} \otimes_{B^{coH}} B})$, constructed in Proposition 2.14, is invariant for the ω -deformation functor, i.e.,*

$$D((M^{coH} \otimes_{B^{coH}} B, \phi_{M^{coH} \otimes_{B^{coH}} B}, \rho_{M^{coH} \otimes_{B^{coH}} B})) = (M^{coH} \otimes_{B^{coH}} B, \phi_{M^{coH} \otimes_{B^{coH}} B}, \rho_{M^{coH} \otimes_{B^{coH}} B}).$$

Proof. To prove the proposition we only need to show that

$$\phi_{M^{coH} \otimes_{B^{coH}} B}^{\omega_{M^{coH} \otimes_{B^{coH}} B}} = \phi_{M^{coH} \otimes_{B^{coH}} B}. \quad (52)$$

Indeed, composing with the coequalizer $n_{M^{coH}} \otimes B$ we have

$$\begin{aligned}
& \phi_{M^{coH} \otimes_{B^{coH}} B}^{\omega_{M^{coH} \otimes_{B^{coH}} B}} \circ (n_{M^{coH}} \otimes B) \stackrel{(43), (42)}{=} n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ (q_B \otimes (\mu_B \circ (h \otimes B))) \circ (\rho_B \otimes B))) \\
& \stackrel{(39)}{=} n_{M^{coH}} \circ (M^{coH} \otimes (\mu_B \circ ((\mu_B \circ (q_B \otimes h)) \circ \rho_B) \otimes B)) \stackrel{(19)}{=} n_{M^{coH}} \circ (M^{coH} \otimes \mu_B) \stackrel{(42)}{=} \phi_{M^{coH} \otimes_{B^{coH}} B} \circ (n_{M^{coH}} \otimes B).
\end{aligned}$$

Therefore, (52) holds. $\square$

The following result is the nonassociative general version of the Fundamental Theorem of Hopf Modules. The proof follows by the properties of the morphism ω_M , obtained in Proposition 2.14, and by (52).

Theorem 2.22. (Fundamental Theorem of Hopf modules) *Assume that the conditions of Proposition 2.14 hold. Let (M, ϕ_M, ρ_M) be an object in $\mathcal{SM}_B^H(h)$. The objects $(M^{coH} \otimes_{B^{coH}} B, \phi_{M^{coH} \otimes_{B^{coH}} B}, \rho_{M^{coH} \otimes_{B^{coH}} B})$ and (M, ϕ_M, ρ_M) are isomorphic in $\mathcal{SM}_B^H(h)$.*

The previous theorem is a generalization of the one proved by Sweedler [30] for Hopf modules over an ordinary Hopf algebra. It also contains the Fundamental Theorem of relative Hopf modules (or (H, B) -Hopf modules, or Doi-Hopf modules) given by Doi and Takeuchi in [14]. On the other hand, in the weak setting, Theorem 2.22 is a generalization of the one obtained by Böhm, Nill and Szlachányi [8], for Hopf modules over a weak Hopf algebra H , and the one proved by Zhang and Zhu [33], for (H, B) -Hopf

modules associated to a weak right H -comodule algebra B . Moreover, in the nonassociative context, it generalizes the result obtained by Brzeziński [10] for Hopf modules associated to a Hopf quasigroup. Finally, for weak Hopf quasigroups, Theorem 2.22 is a generalization of the Fundamental Theorem of Hopf modules proved in [3] (see also [6]).

3. CATEGORICAL EQUIVALENCES FOR STRONG (H, B, h) -HOPF MODULES

As for prerequisites, in this section we will assume that the conditions of Proposition 2.14 hold. Then, in the following H is a weak Hopf quasigroup and (B, ρ_B) is a right H -comodule magma. Also, $h : H \rightarrow B$ is an anchor morphism such that (20) and (39) hold. Finally, the category $\mathcal{C}$ admits coequalizers and the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers. With $\mathcal{C}_{B^{coH}}$ we will denote the category of right B^{coH} -modules.

The main target of this section is to prove that there exists an equivalence between $\mathcal{C}_{B^{coH}}$ and the category of strong (H, B, h) -Hopf modules.

Let (N, ψ_N) be an object in $\mathcal{C}_{B^{coH}}$ and consider the coequalizer diagram

$$\begin{array}{ccc} N \otimes B^{coH} \otimes B & \xrightarrow[\quad N \otimes (\mu_B \circ (i_B \otimes B)) \quad]{\psi_N \otimes B} & N \otimes B \\ & \xrightarrow{n_N} & N \otimes_{B^{coH}} B. \end{array} \quad (53)$$

Then,

$$(n_N \otimes B) \circ (\psi_N \otimes \rho_B) \stackrel{(53)}{=} (n_N \otimes B) \circ (N \otimes ((\mu_B \otimes B) \circ (i_B \otimes \rho_B))) \stackrel{(28)}{=} (n_N \otimes B) \circ (N \otimes (\rho_B \circ (\mu_B \circ (i_B \otimes B))))$$

and, as a consequence, there exists a unique morphism $\rho_{N \otimes_{B^{coH}} B} : N \otimes_{B^{coH}} B \rightarrow N \otimes_{B^{coH}} B \otimes H$ such that

$$\rho_{N \otimes_{B^{coH}} B} \circ n_N = (n_N \otimes B) \circ (N \otimes \rho_B). \quad (54)$$

On the other hand, by (39), we have

$$n_N \circ (\psi_N \otimes \mu_B) = n_N \circ (N \otimes (\mu_B \circ (i_B \otimes \mu_B))) = n_N \circ (N \otimes (\mu_B \circ ((\mu_B \circ (i_B \otimes B)) \otimes B))),$$

and then, using that the functor $- \otimes B$ preserves coequalizers, there exists a unique morphism

$$\phi_{N \otimes_{B^{coH}} B} : N \otimes_{B^{coH}} B \otimes B \rightarrow N \otimes_{B^{coH}} B$$

such that

$$\phi_{N \otimes_{B^{coH}} B} \circ (n_N \otimes B) = n_N \circ (N \otimes \mu_B). \quad (55)$$

By a similar proof to the one used for $M^{coH} \otimes_{B^{coH}} B$ in Proposition 2.14, we can prove that

$$(N \otimes_{B^{coH}} B, \phi_{N \otimes_{B^{coH}} B}, \rho_{N \otimes_{B^{coH}} B})$$

is a strong (H, B, h) -Hopf module such that (the proof follows the ideas given in Proposition 2.21 for $M^{coH} \otimes_{B^{coH}} B$)

$$\phi_{N \otimes_{B^{coH}} B}^{\omega_{N \otimes_{B^{coH}} B}} = \phi_{N \otimes_{B^{coH}} B}. \quad (56)$$

On the other hand, if $f : N \rightarrow P$ is a morphism in $\mathcal{C}_{B^{coH}}$, we have

$$n_P \circ (f \otimes B) \circ (\psi_N \otimes B) = n_P \circ (f \otimes B) \circ (N \otimes (\mu_B \circ (i_B \otimes B)))$$

and then there exists a unique morphism $f \otimes_{B^{coH}} B : N \otimes_{B^{coH}} B \rightarrow P \otimes_{B^{coH}} B$ such that

$$n_P \circ (f \otimes B) = (f \otimes_{B^{coH}} B) \circ n_N. \quad (57)$$

The morphism $f \otimes_{B^{coH}} B$ is a morphism in $\mathcal{SM}_B^H(h)$ because

$$\begin{aligned} \rho_{P \otimes_{B^{coH}} B} \circ f \otimes_{B^{coH}} B \circ n_N &\stackrel{(54), (57)}{=} (n_P \otimes B) \circ (f \otimes \rho_B) \stackrel{(54)}{=} \rho_{P \otimes_{B^{coH}} B} \circ n_P \circ (f \otimes B) \\ &\stackrel{(57)}{=} (f \otimes_{B^{coH}} B \otimes B) \circ \rho_{N \otimes_{B^{coH}} B} \circ n_N \end{aligned}$$

and

$$\phi_{P \otimes_{B^{coH}} B}^{\omega_{P \otimes_{B^{coH}} B}} \circ (f \otimes_{B^{coH}} B \otimes B) \circ (n_N \otimes B) \stackrel{(56)}{=} \phi_{P \otimes_{B^{coH}} B} \circ (f \otimes_{B^{coH}} B \otimes B) \circ (n_N \otimes B) \stackrel{(55), (57)}{=} n_P \circ (f \otimes \mu_B)$$

$$\stackrel{(57)}{=} f \otimes_{B^{coH}} B \circ n_N \circ (N \otimes \mu_B) \stackrel{(55)}{=} f \otimes_{B^{coH}} B \circ \phi_{N \otimes_{B^{coH}} B} \circ (n_N \otimes B) \stackrel{(56)}{=} f \otimes_{B^{coH}} B \circ \phi_{N \otimes_{B^{coH}} B}^{\omega_N \otimes_{B^{coH}} B} \circ (n_N \otimes B).$$

Summarizing, we have the following proposition:

Proposition 3.1. *There exists a functor $F : \mathcal{C}_{B^{coH}} \rightarrow \mathcal{SM}_B^H(h)$, called the induction functor, defined on objects by $F((N, \psi_N)) = (N \otimes_{B^{coH}} B, \phi_{N \otimes_{B^{coH}} B}, \rho_{N \otimes_{B^{coH}} B})$ and on morphisms by $F(f) = f \otimes_{B^{coH}} B$.*

Now let (M, ϕ_M, ρ_M) be an object in $\mathcal{SM}_B^H(h)$. By Proposition 2.13 we have that the object of coinvariants M^{coH} is a right B^{coH} -module where $\phi_{M^{coH}} = p_M \circ \phi_M \circ (i_M \otimes i_B)$ (see (38)). Let $g : M \rightarrow Q$ be a morphism in $\mathcal{SM}_B^H(h)$. Using the comodule morphism condition we obtain that $\rho_Q \circ g \circ i_M = (Q \otimes \overline{\Pi}_H^R) \circ \rho_Q \circ g \circ i_M$ and this implies that there exists a unique morphism $g^{coH} : M^{coH} \rightarrow Q^{coH}$ such that

$$i_Q \circ g^{coH} = g \circ i_M. \quad (58)$$

Also, using that g is B -quasilinear, H colinear, and (49) we have

$$g \circ q_M^{\omega_M} = g \circ q_M = q_T \circ g = q_T^{\alpha_T} \circ g. \quad (59)$$

Then,

$$i_Q \circ g^{coH} \circ p_M \stackrel{(58)}{=} g \circ q_M \stackrel{(59)}{=} q_Q \circ g$$

and, as a consequence,

$$g^{coH} \circ p_M = p_Q \circ g \quad (60)$$

holds.

The morphism g^{coH} is a morphism in $\mathcal{C}_{B^{coH}}$ because

$$\begin{aligned} & \phi_{Q^{coH}} \circ (g^{coH} \otimes B^{coH}) \\ &= p_Q \circ \phi_Q \circ ((i_Q \circ g^{coH}) \otimes i_B) \quad ((38)) \\ &= p_Q \circ \phi_Q^{\omega_Q} \circ ((i_Q \circ g^{coH}) \otimes i_B) \quad ((51)) \\ &= p_Q \circ \phi_Q^{\omega_Q} \circ ((g \circ i_M) \otimes i_B) \quad ((58)) \\ &= p_Q \circ g \circ \phi_M^{\omega_M} \circ (i_M \otimes i_B) \quad ((45)) \\ &= p_Q \circ g \circ \phi_M \circ (i_M \otimes i_B) \quad ((51)) \\ &= g^{coH} \circ \phi_{M^{coH}} \quad ((60)). \end{aligned}$$

Thus, we have the following result.

Proposition 3.2. *There exists a functor $G : \mathcal{SM}_B^H(h) \rightarrow \mathcal{C}_{B^{coH}}$, called the functor of coinvariants, defined on objects by $G((M, \phi_M, \rho_M)) = (M^{coH}, \psi_{M^{coH}})$ and on morphisms by $G(g) = g^{coH}$.*

Theorem 3.3. *The categories $\mathcal{SM}_B^H(h)$ and $\mathcal{C}_{B^{coH}}$ are equivalent.*

Proof. To prove the theorem, we firstly obtain that the induction functor F , introduced in Proposition 3.1, is left adjoint to the functor of coinvariants G introduced in Proposition 3.2. Later, we show that the unit and counit associated to this adjunction are natural isomorphisms. Then, we proceed as in the proof of Theorem 3.10 of [6] by dividing the proof in three steps.

Step 1: In this step we define the unit of the adjunction. For any right B^{coH} -module (N, ψ_N) , consider

$$\alpha_N : N \rightarrow GF(N) = (N \otimes_{B^{coH}} B)^{coH}$$

as the unique morphism such that

$$i_{N \otimes_{B^{coH}} B} \circ \alpha_N = n_N \circ (N \otimes \eta_B). \quad (61)$$

This morphism exists and is unique because

$$\begin{aligned} & ((N \otimes_{B^{coH}} B) \otimes \overline{\Pi}_H^R) \circ \rho_{N \otimes_{B^{coH}} B} \circ n_N \circ (N \otimes \eta_B) \stackrel{(54)}{=} (n_N \otimes H) \circ (N \otimes ((B \otimes \overline{\Pi}_H^R) \circ \rho_B \circ \eta_B)) \\ & \stackrel{(b5)}{=} (n_N \otimes H) \circ (N \otimes (\rho_B \circ \eta_B)) \stackrel{(54)}{=} \rho_{N \otimes_{B^{coH}} B} \circ n_N \circ (N \otimes \eta_B). \end{aligned}$$

Also, α_N is a morphism in $\mathcal{C}_{B^{coH}}$. Indeed: Composing with the equalizer $i_{N \otimes_{B^{coH}} B}$ we have

$$i_{N \otimes_{B^{coH}} B} \circ \psi_{N \otimes_{B^{coH}} B} \circ (\alpha_N \otimes B^{coH})$$

$$\begin{aligned}
&= q_{N \otimes_{B^{coH}} B} \circ \phi_{N \otimes_{B^{coH}} B} \circ ((i_{N \otimes_{B^{coH}} B} \circ \alpha_N) \otimes i_B) \quad ((38)) \\
&= q_{N \otimes_{B^{coH}} B} \circ \phi_{N \otimes_{B^{coH}} B} \circ ((n_N \circ (N \otimes \eta_B)) \otimes i_B) \quad ((61)) \\
&= q_{N \otimes_{B^{coH}} B} \circ n_N \circ (N \otimes (\mu_B \circ (\eta_B \otimes i_B))) \quad ((55)) \\
&= n_N \circ (N \otimes (q_B \circ i_B)) \quad (\text{properties of } \eta_B, (54), \text{ and } (55)) \\
&= n_N \circ (N \otimes i_B) \quad (\text{properties of } i_B) \\
&= n_N \circ (N \otimes (\mu_B \circ (i_B \otimes \eta_B))) \quad (\text{properties of } \eta_B) \\
&= n_N \circ (\psi_N \otimes \eta_B) \quad ((53)) \\
&= i_{N \otimes_{B^{coH}} B} \circ \alpha_N \circ \psi_N \quad ((61))
\end{aligned}$$

and therefore $\psi_{N \otimes_{B^{coH}} B} \circ (\alpha_N \otimes B^{coH}) = \alpha_N \circ \psi_N$. On the other hand, the morphism α_N is natural in N because if $f : N \rightarrow P$ is a morphism in $\mathcal{C}_{B^{coH}}$

$$\begin{aligned}
i_{P \otimes_{B^{coH}} B} \circ (f \otimes_{B^{coH}} B)^{coH} \circ \alpha_N &\stackrel{(58)}{=} (f \otimes_{B^{coH}} B) \circ i_{N \otimes_{B^{coH}} B} \circ \alpha_N \stackrel{(61)}{=} (f \otimes_{B^{coH}} B) \circ n_N \circ (N \otimes \eta_B) \\
&\stackrel{(57)}{=} n_P \circ (f \otimes \eta_B) \stackrel{(61)}{=} i_{P \otimes_{B^{coH}} B} \circ \alpha_P \circ f,
\end{aligned}$$

and then $(f \otimes_{B^{coH}} B)^{coH} \circ \alpha_N = \alpha_P \circ f$.

Finally, we prove that α_N is an isomorphism for all right B^{coH} -module N . First note that, under the conditions of this theorem, the triple (B, μ_B, δ_B) is a strong (H, B, h) -Hopf module, and then

$$\psi_N \circ (\psi_N \otimes p_B) = \psi_N \circ (N \otimes (p_B \circ \mu_B \circ (i_B \otimes B)))$$

because

$$\begin{aligned}
&\psi_N \circ (\psi_N \otimes p_B) \\
&= \psi_N \circ (N \otimes (\mu_{B^{coH}} \circ (B^{coH} \otimes p_B))) \quad (\text{module condition for } N) \\
&= \psi_N \circ (N \otimes (p_B \circ \mu_B \circ (i_B \otimes q_B))) \quad ((12)) \\
&= \psi_N \circ (N \otimes (p_B \circ \mu_B \circ (i_B \otimes B))) \quad ((33)).
\end{aligned}$$

Therefore, there exists a unique morphism $m_N : N \otimes_{B^{coH}} B \rightarrow N$ such that

$$m_N \circ n_N = \psi_N \circ (N \otimes p_B). \quad (62)$$

Now define the morphism $x_N : (N \otimes_{B^{coH}} B)^{coH} \rightarrow N$ by $x_N = m_N \circ i_{N \otimes_{B^{coH}} B}$. Then,

$$x_N \circ \alpha_N \stackrel{(61)}{=} m_N \circ n_N \circ (N \otimes \eta_B) \stackrel{(62)}{=} \psi_N \circ (N \otimes (p_B \circ \eta_B)) \stackrel{(9)}{=} \psi_N \circ (N \otimes \eta_{B^{coH}}) = id_N.$$

On the other hand, composing with $i_{N \otimes_{B^{coH}} B}$ and $p_{N \otimes_{B^{coH}} B} \circ n_N$ we have

$$\begin{aligned}
&i_{N \otimes_{B^{coH}} B} \circ \alpha_N \circ x_N \circ p_{N \otimes_{B^{coH}} B} \circ n_N \\
&= n_N \circ ((m_N \circ q_{N \otimes_{B^{coH}} B} \circ n_N) \otimes \eta_B) \quad ((61)) \\
&= n_N \circ ((\psi_N \circ (N \otimes (p_B \circ q_B))) \otimes \eta_B) \quad ((54), (55), (62)) \\
&= n_N \circ ((\psi_N \circ (N \otimes p_B)) \otimes \eta_B) \quad (\text{properties of } q_B) \\
&= n_N \circ (N \otimes (\mu_B \circ (q_B \otimes \eta_B))) \quad ((53)) \\
&= n_N \circ (N \otimes q_B) \quad (\text{properties of } \eta_B) \\
&= q_{N \otimes_{B^{coH}} B} \circ n_N \quad ((54), (55)).
\end{aligned}$$

Therefore, $\alpha_N \circ x_N = id_{(N \otimes_{B^{coH}} B)^{coH}}$ and, as a consequence, α_N is an isomorphism.

Step 2: For any $(M, \phi_M, \rho_M) \in \mathcal{SM}_B^H(h)$ the counit is defined as $\beta_M = \omega_M : M^{coH} \otimes_{B^{coH}} B \rightarrow M$, where ω_M is the isomorphism satisfying (44). By Theorem 2.22, we know that ω_M is an isomorphism in $\mathcal{SM}_B^H(h)$, and it is natural because if $g : M \rightarrow Q$ is a morphism in $\mathcal{SM}_B^H(h)$ we have

$$\begin{aligned}
\beta_Q \circ (g^{coH} \otimes_{B^{coH}} B) \circ n_{M^{coH}} &\stackrel{(57)}{=} \beta_Q \circ n_{Q^{coH}} \circ (g^{coH} \otimes B) \stackrel{(44)}{=} \phi_Q \circ ((i_Q \circ g^{coH}) \otimes B) \stackrel{(51)}{=} \phi_Q^{\omega_Q} \circ ((i_Q \circ g^{coH}) \otimes B) \\
&\stackrel{(58)}{=} \phi_Q^{\omega_Q} \circ ((g \circ i_Q) \otimes B) \stackrel{(45)}{=} g \circ \phi_M^{\omega_M} \circ (i_M \otimes B) \stackrel{(51)}{=} g \circ \phi_M \circ (i_M \otimes B) \stackrel{(44)}{=} g \circ \beta_M \circ n_{M^{coH}}
\end{aligned}$$

and thus $\beta_Q \circ g^{coH} \otimes_{B^{coH}} B = g \circ \beta_M$.

Step 3: Now we prove the triangular identities for the unit and counit previously defined. Indeed: The first triangular identity holds because composing with n_N we have

$$\beta_{N \otimes_{B^{coH}} B} \circ (\alpha_N \otimes_{B^{coH}} B) \circ n_N$$

$$\begin{aligned}
&= \beta_{N \otimes_{B^{coH}} B} \circ n_{(N \otimes_{B^{coH}} B)^{coH}} \circ (\alpha_N \otimes B) \text{ (by (57))} \\
&= \phi_{N \otimes_{B^{coH}} B} \circ ((i_{N \otimes_{B^{coH}} B} \circ \alpha_N) \otimes B) \text{ (by (44))} \\
&= \phi_{N \otimes_{B^{coH}} B} \circ ((n_N \circ (N \otimes \eta_B)) \otimes B) \text{ (by (61))} \\
&= n_N \circ (N \otimes (\mu_B \circ (\eta_B \otimes B))) \text{ (by (55))} \\
&= n_N \text{ (by the unit properties).}
\end{aligned}$$

Finally, if we compose with i_M ,

$$i_M \circ \beta_M^{coH} \circ \alpha_{M^{coH}} \stackrel{(58)}{=} \beta_M \circ i_{N \otimes_{B^{coH}} B} \circ \alpha_{M^{coH}} \stackrel{(61)}{=} \beta_M \circ n_{M^{coH}} \circ (M^{coH} \otimes \eta_B) \stackrel{(44)}{=} \phi_M \circ (i_M \otimes \eta_B) \stackrel{(d2-1)}{=} i_M,$$

and then $\beta_M^{coH} \circ \alpha_{M^{coH}} = id_{M^{coH}}$.

□

As a consequence, we obtain the following particular instances of our main theorem.

Corollary 3.4. *The following assertions hold:*

- (i) *Let H be a Hopf algebra and let (B, ρ_B) be a right H -comodule monoid such the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers. Then, if there exists a multiplicative total integral $h : H \rightarrow B$, the categories of right (H, B) -Hopf modules, denoted by $\mathcal{M}_B^H$ and introduced by Doi in [13], the category $\mathcal{SM}_B^H(h)$ of strong (H, B, h) -Hopf modules, and the category of right B^{coH} -modules $\mathcal{C}_{B^{coH}}$ are equivalent. In particular, if $B = H$ and $\rho_B = \delta_H$, the Sweedler category of Hopf modules $\mathcal{M}_H^H$, the category $\mathcal{SM}_H^H(id_H)$ of strong (H, H, id_H) -Hopf modules, and the category $\mathcal{C}$ are equivalent.*
- (ii) *Let H be a weak Hopf algebra and let (B, ρ_B) be a right H -comodule monoid such that the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers. Then, if there exists a multiplicative total integral $h : H \rightarrow B$, the categories of right (H, B) -Hopf modules, denoted by $\mathcal{M}_B^H$ and introduced by Böhm in [9] (see also [33] and [16] for the categorical equivalence), the category $\mathcal{SM}_B^H(h)$ of strong (H, B, h) -Hopf modules, and the category of right B^{coH} -modules $\mathcal{C}_{B^{coH}}$ are equivalent. In particular, if $B = H$ and $\rho_B = \delta_H$, the category of Hopf modules $\mathcal{M}_H^H$, the category $\mathcal{SM}_H^H(id_H)$ of strong (H, H, id_H) -Hopf modules, and the category $\mathcal{C}_{H_L}$ of right H_L -modules are equivalent.*
- (iii) *Let H be a Hopf quasigroup and let (B, ρ_B) be a right H -comodule magma such the functors $- \otimes B$ and $- \otimes H$ preserve coequalizers. Then, if there exists an anchor morphism $h : H \rightarrow B$, the categories $\mathcal{SM}_B^H(h)$ of strong (H, B, h) -Hopf modules, and the category of right B^{coH} -modules $\mathcal{C}_{B^{coH}}$ are equivalent. In particular, if $B = H$ and $\rho_B = \delta_H$, we obtain the result proved by Brzeziński in [10]: the category of Hopf modules $\mathcal{M}_H^H = \mathcal{SM}_H^H(id_H)$ and the category $\mathcal{C}$ are equivalent.*
- (iv) *Let H be a weak Hopf quasigroup such the functor $- \otimes H$ preserves coequalizers. The category of strong Hopf modules $\mathcal{SM}_H^H = \mathcal{SM}_H^H(id_H)$, and the category $\mathcal{C}_{H_L}$ of right H_L -modules are equivalent (this is the main result proved in [6]).*

ACKNOWLEDGEMENTS

The authors were supported by Ministerio de Economía y Competitividad (Spain), grant MTM2016-79661-P. AEI/FEDER, UE, support included (Homología, homotopía e invariantes categóricos en grupos y álgebras no asociativas).

REFERENCES

- [1] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R, Soneira Calvo, C. Projections and Yetter-Drinfel'd modules over Hopf (co)quasigroups. J. Algebra, 2015, 443: 153-199.
- [2] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R, Soneira Calvo, C. Cleft comodules over Hopf quasigroups. Commun. Contemp. Math., 2015, 17: 1550007.
- [3] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R. Weak Hopf quasigroups. Asian J. of Math., 2016, 20: 665-694.
- [4] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R. Cleft and Galois extensions associated to a weak Hopf quasigroup. J. Pure Appl. Algebra, 2016, 220: 1002-1034.

- [5] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R. A characterization of weak Hopf (co)quasigroups. *Mediterr. J. Math.*, 2016, 13: 3747-3764.
- [6] Alonso Álvarez J N, Fernández Vilaboa J M, González Rodríguez R. Strong Hopf modules for weak Hopf quasigroups. *Colloq. Math. Warsaw* (2017) (in press) (available in arXiv:1505.04586).
- [7] Benkart G., Madariaga S, Pérez-Izquierdo, J M. Hopf algebras with triality. *Trans. Amer. Math. Soc.*, 2012, 365: 1001-1023.
- [8] Böhm G, Nill F, Szlachányi K. Weak Hopf algebras, I. Integral theory and C^* -structure. *J. Algebra*, 1999, 221: 385-438.
- [9] Böhm G. Doi-Hopf modules over weak Hopf algebras. *Comm. Algebra*, 2000, 10: 4687-4698.
- [10] Brzeziński T. Hopf modules and the fundamental theorem for Hopf (co)quasigroups. *Internat. Elec. J. Algebra*, 2010, 8: 114-128.
- [11] Brzeziński T, Jiao Z. R-smash products of Hopf quasigroups. *Arab. J. Math.*, 2012, 1: 39-46.
- [12] Brzeziński T, Jiao Z. Actions of Hopf quasigroups. *Comm. Algebra*, 2012, 40: 681-696.
- [13] Doi Y. On the structure of relative Hopf modules. *Comm. Algebra*, 1983, 11: 243-255.
- [14] Doi Y, Takeuchi M. Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and Azumaya algebras, *J. Algebra*, 1989, 121: 488-516.
- [15] Fang X, Torrecillas B. Twisted smash products and L-R smash products for biquasimodule Hopf quasigroups. *Comm. Algebra*, 2014, 42:4204-4234.
- [16] Henker H. Module categories over quasi-Hopf algebras and weak Hopf algebras and the projectivity of Hopf modules. Dissertation for the Doctoral Degree. Munich: LMU Munich, 2011 (available in <http://edoc.ub.uni-muenchen.de/13148/>)
- [17] Klim J. Bicrossproduct Hopf quasigroups. *Comment. Math. Univ. Carolinae*, 2010, 51: 287-304.
- [18] Jiao Z, Meng H. Ore extensions of Hopf coquasigroups, *Math. Notes*, 2014, 95: 338-345.
- [19] Jiao Z, Zhao X. Almost cocommutative and quasitriangular Hopf coquasigroups. *J. Algebra Appl.*, 2014, 13: 1450010.
- [20] Klim J, Majid S. Hopf quasigroups and the algebraic 7-sphere. *J. Algebra*, 2010, 323: 3067-3110.
- [21] Larson R G, Sweedler M E. An associative orthogonal bilinear form for Hopf algebras. *Amer. J. Math.*, 1969, 91: 75-93.
- [22] Mac Lane S. *Categories for the Working Mathematician*. GTM 5, New York: Springer-Verlag, New-York, 1971.
- [23] Mostovoy J, Pérez-Izquierdo J M, Shestakov I P. Hopf algebras in non associative Lie theory. *Bull. Math. Sci.*, 2014, 4: 129-173.
- [24] Pérez-Izquierdo J M, Shestakov I P. An envelope for Malcev algebras. *J. Algebra*, 2004, 272: 379-393.
- [25] Pérez-Izquierdo J M. Algebras, hyperalgebras, nonassociative bialgebras and loops. *Adv. Math.*, 2007, 208: 834-876.
- [26] Smith J D H. One-sided Quantum quasigroups and loops. *Demonstr. Math.*, 2015, 48: 620-636.
- [27] Smith J D H. Quantum quasigroups and loops. *J. Algebra*, 2016, 456: 46-75.
- [28] Smith J D H. Quantum idempotence, distributivity, and the Yang-Baxter equation. *Comment. Math. Univ. Carolin.*, 2016, 57: 567-583.
- [29] Smith J D H. Quantum Quasigroups and the Quantum Yang-Baxter Equation. *Axioms*, 2016, 5: 25.
- [30] Sweedler M E. *Hopf algebras*. New York: Benjamin, New York 1969.
- [31] Tvalavadze M V, Bremner M R. Enveloping algebras of solvable Malcev algebras of dimension five. *Comm. in Algebra*, 2011, 39-1532-4125.
- [32] Takeuchi M. Relative Hopf-modules-equivalences and freeness criteria. *J. Algebra*, 1979, 60: 452-471.
- [33] Zhang L, Zhu S. Fundamental theorems of weak Doi-Hopf modules and semisimple weak smash product Hopf algebras. *Comm. Algebra*, 2004, 32:3403-3415.